

PATENT

ATTORNEY DOCKET NO. CGNE.115.01US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: McBride *et al* ) Examiner: Campell, B.  
Serial No.: 08/984,099 )  
Filed: December 3, 1997 ) Art Unit: 1632  
Title: COTTON FIBER TRANSCRIPTIONAL )  
FACTORS )  
COMMUNICATION

Bruce Campell  
c/o Technology Center 1600  
Crystal Mall 1, 7<sup>th</sup> Floor  
1911 South Clark St.  
Arlington, VA 22202

Sir:

As per our facsimile to you of December 15, 1999, please find enclosed two color drawings and a copy of PCT 96/09897 in the above-entitled patent application.

Respectfully submitted,

Date: December 17, 1999

Barbara Rae-Venter

Barbara Rae-Venter, Ph.D.  
Reg. No. 32,750

Rae-Venter Law Group, P.C.  
P. O. Box 60039  
Palo Alto, CA 94306  
Telephone: (650) 328-4400  
Facsimile: (650) 328-4477  
BRV/JLW

Hand Delivery  
CERTIFICATE OF EXPRESS MAILING

"Express Mail" Label No.

Date of Deposit: 12/20/99

I hereby certify under 37 C.F.R. 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to the Commissioner of Patents and Trademarks, Washington, D.C., 20231.

(Signature)

JAMES W. BURTON  
(Printed Name)

RECEIVED  
TECH. CENTER 1600  
- 99 DEC 20  
PM 12  
200

08/984, 099

Attachment #12

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/29, 15/82, 5/10, A01H 5/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 96/40924</b>
			(43) International Publication Date: 19 December 1995 (19.12.95)
(21) International Application Number: PCT/US96/09897		(81) Designated States: AU, CA, CN, JP, KG, KZ, MX, TJ, TM, TR, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 7 June 1995 (07.06.95)			
(30) Priority Data: 08/480,178 7 June 1995 (07.06.95) US		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(60) Parent Application or Grant (63) Related by Continuation US Filed on		Not furnished (CIP) Not furnished	
(71) Applicant (for all designated States except US): CALGENE, INC. [US/US]; 1920 Fifth Street, Davis, CA 95616 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): McBRIDE, Kevin [US/US]; 1309 Marina Circle, Davis, CA 95616 (US). STALKER, David, M. [US/US]; 2736 Cumberland Place, Davis, CA 95616 (US). PEAR, Julie, R. [US/US]; 818 Douglas Avenue, Davis, CA 95616 (US). PEREZ-GRAU, Luis [ES/US]; 1230 Elk Place, Davis, CA 95616 (US).			
(74) Agents: SCHWEDLER, Carl, J. et al.; Calgene, Inc., 1920 Fifth Street, Davis, CA 95616 (US).			
(54) Title: COTTON FIBER TRANSCRIPTIONAL FACTORS			
(57) Abstract  Novel DNA constructs are provided which may be used as molecular probes or inserted into a plant host to provide for modification of transcription of a DNA sequence of interest during various stages of cotton fiber development. The DNA constructs comprise a cotton fiber transcriptional initiation regulatory region associated with a gene which is expressed in cotton fiber. Also provided is novel cotton having a cotton fiber which has a natural color introduced by the expression in the cotton fiber cell, using such a construct, of pigment synthesis genes. Cotton fiber cells having color produced by genetic engineering and cotton cells comprising melanin and indigo pigments are included.			

No 371 filed

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## COTTON FIBER TRANSCRIPTIONAL FACTORS

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation in part of United States application Serial No. 08/487,087 filed June 7, 1995, and a continuation in part of United States application Serial No. 08/480,178, filed June 7, 1995.

INTRODUCTIONTechnical Field

10 This invention relates to methods of using *in vitro* constructed DNA transcription or expression cassettes capable of directing fiber-tissue transcription of a DNA sequence of interest  
15 in plants to produce fiber cells having an altered phenotype, and to methods of providing for or modifying various characteristics of cotton fiber. The invention is exemplified by methods of using cotton fiber promoters for altering the phenotype of cotton fiber, and cotton fibers produced by the method.

Background

20 In general, genetic engineering techniques have been directed to modifying the phenotype of individual prokaryotic and eukaryotic cells, especially in culture. Plant cells have proven  
25 more intransigent than other eukaryotic cells, due not only to a lack of suitable vector systems but also as a result of the different goals involved. For many applications, it is desirable

to be able to control gene expression at a particular stage in the growth of a plant or in a particular plant part. For this purpose, regulatory sequences are required which afford the desired initiation of transcription in the appropriate cell types and/or at the appropriate time in the plant's development without having serious detrimental effects on plant development and productivity. It is therefore of interest to be able to isolate sequences which can be used to provide the desired regulation of transcription in a plant cell during the growing cycle of the host plant.

One aspect of this interest is the ability to change the phenotype of particular cell types, such as differentiated epidermal cells that originate in fiber tissue, i.e. cotton fiber cells, so as to provide for altered or improved aspects of the mature cell type. Cotton is a plant of great commercial significance. In addition to the use of cotton fiber in the production of textiles, other uses of cotton include food preparation with cotton seed oil and animal feed derived from cotton seed husks.

Despite the importance of cotton as a crop, the breeding and genetic engineering of cotton fiber phenotypes has taken place at a relatively slow rate because of the absence of reliable promoters for use in selectively effecting changes in the phenotype of the fiber. In order to effect the desired phenotypic changes, transcription initiation regions capable of initiating transcription in fiber cells during development are desired. Thus, an important goal of cotton bioengineering research is the

acquisition of a reliable promoter which would permit expression of a protein selectively in cotton fiber to affect such qualities as fiber strength, length, color and dyability.

5 Relevant Literature

Cotton fiber-specific promoters are discussed in PCT publications WO 94/12014 and WO 95/08914, and John and Crow, Proc. Natl. Acad. Sci. USA, 89:5769-5773, 1992. cDNA clones that are preferentially expressed in cotton fiber have been isolated. One  
10 of the clones isolated corresponds to mRNA and protein that are highest during the late primary cell wall and early secondary cell wall synthesis stages. John and Crow, *supra*.

In animals, the *ras* superfamily is subdivided into the subfamilies *ras* which is involved in controlling cell growth and  
15 division, *rab/YPT* members which control secretory processes, and *rho* which is involved in control of cytoskeletal organization (Bourne et al., (1991) Nature 349: 117-127), and number of homologous genes have now been identified in plants (for a review, see Terryn et al., (1993) Plant Mol. Biol. 22: 143-152). None  
20 have been found for the important *ras* subfamily, all but one of the genes identified have been members of the *rab/YPT1* subfamily, and there is only one recent report of the cloning of a *rho* gene in pea (Yang and Watson(1993) Proc. Natl. Acad. Sci. USA 90: 8732-8736).

25 Little work has been done to characterize the functions of these genes in plants, although one recent report has shown that a small G protein from *Arabidopsis* can functionally complement a

mutant form in yeast involved in vesicle trafficking, suggesting a similar function for the plant gene (Bednarek et al., (1994) Plant Physiol 104: 591-596).

In animals, two members of the *rho* subfamily, called Rac and Rho, have been shown to be involved in the regulation of actin organization (for a review, see Downward, (1992) Nature 359: 273-274).

Rac1 has been shown to mediate growth factor-induced membrane ruffling by influencing microfilament alignment on the plasma membrane (Ridley et al, (1992) Cell 70: 401-410), whereas RhoA regulates the formation of actin stress fibers associated with focal adhesions (Ridley and Hall, (1992) Cell 70: 389-399).

In yeast, the CDC42 gene codes for a *rho*-type protein which also regulates actin organization involved in the establishment of cell polarity required for the localized deposition of chitin in the bud scar (Adams et al., (1990) J Cell Biol 111: 131-143).

Disruption of gene function, either by temperature shifts with a CDC42-temperature-sensitive mutant in yeast (Adams et al., 1990), or by micro-injection into fibroblasts of mutant Rac or Rho proteins exhibiting a dominant negative phenotype (Ridley et al., 1992; Ridley and Hall, 1992), leads to disorganization of the actin network.

In plants, control of cytoskeletal organization is poorly understood in spite of its importance for the regulation of patterns of cell division, expansion, and subsequent deposition of secondary cell wall polymers. The cotton fiber represents an excellent system for studying cytoskeletal organization. Cotton

fibers are single cells in which cell elongation and secondary wall deposition can be studied as distinct events. These fibers develop synchronously within the boll following anthesis, and each fiber cell elongates for about 3 weeks, depositing a thin primary wall (Meinert and Delmer, (1984) Plant Physiol. 59: 1088-1097; Basra and Malik, (1984) Int Rev of Cytol 89: 65-113). At the time of transition to secondary wall cellulose synthesis, the fiber cells undergo a synchronous shift in the pattern of cortical microtubule and cell wall microfibril alignments, events which may be regulated upstream by the organization of actin (Seagull, (1990) Protoplasma 159: 44-59; and (1992) In: Proceedings of the Cotton Fiber Cellulose Conference, National Cotton Council of America, Memphis TN, pp 171-192.

Agrobacterium-mediated cotton transformation is described in Umbeck, United States Patents Nos. 5,004,863 and 5,159,135 and cotton transformation by particle bombardment is reported in WO 92/15675, published September 17, 1992. Transformation of Brassica has been described by Radke et al. (Theor. Appl. Genet. (1988) 75:685-694; Plant Cell Reports (1992) 11:499-505.

#### SUMMARY OF THE INVENTION

Novel DNA constructs and methods for their use are described which are capable of directing transcription of a gene of interest in cotton fiber, particularly early in fiber development and during secondary cell wall development. The novel constructs include a vector comprising a transcriptional and translational initiation region obtainable from a gene expressed in cotton fiber



and methods of using constructs including the vector for altering fiber phenotype. Both the endogenous 3' regions and 5' regions may be important in directing efficient transcription and translation.

5        Three promoters are provided from genes involved in the regulation of cotton fiber development. One, Rac13, is from a protein in cotton which codes for an animal Rac protein homolog. Rac13, shows highly-enhanced expression during fiber development. This pattern of expression correlates well with the timing of  
10        reorganization of the cytoskeleton, suggesting that the Rac13 cotton gene may, like its animal counterpart, be involved in the signal transduction pathway for cytoskeletal organization. Rac13 is a gene that is moderately expressed during fiber development turning on at 9 dpa and shutting down approximately 24 dpa. It is  
15        maximally expressed between 17-21 dpa developing fiber.

Another promoter from a cotton protein is designated 4-4. The 4-4 mRNA accumulates in fiber cells at day 17 post anthesis and continues towards fiber maturity, which occurs at 60 days or so post anthesis. Data demonstrates that the 4-4 promoter remains  
20        very active at day 35 post anthesis.

Also provided is a promoter from a lipid transfer protein (hereinafter sometimes referred to as "Ltp") which is preferentially expressed in cotton fiber.

The methods of the present invention include transfecting a  
25        host plant cell of interest with a transcription or expression cassette comprising a cotton fiber promoter and generating a plant which is grown to produce fiber having the desired phenotype.

Constructs and methods of the subject invention thus find use in modulation of endogenous fiber products, as well as production of exogenous products and in modifying the phenotype of fiber and fiber products. The constructs also find use as molecular probes.

- 5 In particular, constructs and methods for use in gene expression in cotton embryo tissues are considered herein. By these methods, novel cotton plants and cotton plant parts, such as modified cotton fibers, may be obtained.

- Also provided are constructs and methods of use relating to  
10 modification of color phenotype in cotton fiber. Such constructs contain sequences for expression of genes involved in the production of colored compounds, such as anthocyanins, melanin or indigo, and also may contain sequences which provide for targeting of the gene products to particular locations in the plant cell,  
15 such as plastid organelles, or vacuoles. Plastid targeting is of particular interest for expression of genes involved in aromatic amino acid biosynthesis pathways, while vacuolar targeting is of particular interest where the precursors required in synthesis of the pigment are present in vacuoles.

- 20 Of particular interest are plants producing fibers which are color, that is, with pigment produced in the fiber by the plant during fiber development, as opposed to fibers which are harvested and dyed or otherwise pigmented by separate processing. Fibers from a plant producing such colored fiber may be used to produce  
25 colored yarns and/or fabric which have not been subjected to any dyeing process. While naturally colored cotton has been available from various domesticated and wild type cotton varieties, the

instant application provides cotton fiber has a color produced by the expression of a genetically engineered protein.

Thus, the application provides constructs and methods of use relating to modification of color phenotype in cotton fiber. Such  
5 constructs contain sequences for expression of genes involved in the production of colored compounds, such as melanin or indigo, and also contain sequences which provide for targeting of the gene products to particular locations in the plant cell, such as  
10 plastid organelles, or vacuoles. Plastid targeting is of particular interest for expression of genes involved in the aromatic amino acid biosynthesis pathways, while vacuolar targeting is of particular interest where the precursors required in synthesis of the pigment are present in vacuoles.

15

#### DESCRIPTION OF THE DRAWINGS

Figure 1 shows the DNA sequence encoding the structural protein from cDNA 4-4.

Figure 2 shows the sequence to the promoter construct pCGN5606 made using genomic DNA from 4-4-6 genomic clone.

20 Figure 3 shows the sequence to the 4-4 promoter construct pCGN5610.

Figure 4 shows the cDNA sequence encoding the Rac13 gene expressed in cotton fiber.

25 Figure 5 shows the sequence the promoter region from the rac13 gene.

Figure 6 shows a restriction map for pCGN4735.

Figure 7 shows the sequence of the Ltp promoter region from a cotton fiber specific lipid transfer protein gene.

Figure 8 shows the arrangement of a binary vectors pCGN5148 and pCGN5616 for plant transformation to express genes for melanin  
5 synthesis and indigo synthesis, respectively.

Figure 9 provides the results of color measurements taken from fibers of the control Coker 130 cotton used in transformation using color constructs.

Figure 10 shows the results of measurements made of color of  
10 plants transformed by the pCGN5148 construct to express genes for melanin synthesis.

Figure 11 shows the results of measurements taken of the color of plants transformed by the pCGN5149 construct to express genes for melanin synthesis.

15 Figure 12 shows the results of measurements made of color of plants transformed to express genes for indigo synthesis, using construct pCGN5616.

Figure 13 shows control measurements made of naturally colored cotton plants which are produced by non-transgenic colored  
20 cotton plants.

#### DETAILED DESCRIPTION OF THE INVENTION

In accordance with the subject invention, novel constructs and methods are described, which may be used provide for  
25 transcription of a nucleotide sequence of interest in cells of a plant host, preferentially in cotton fiber cells to produce cotton fiber having an altered color phenotype.

Cotton fiber is a differentiated single epidermal cell of the outer integument of the ovule. It has four distinct growth phases; initiation, elongation (primary cell wall synthesis), secondary cell wall synthesis, and maturation. Initiation of  
5 fiber development appears to be triggered by hormones. The primary cell wall is laid down during the elongation phase, lasting up to 25 days postanthesis (DPA). Synthesis of the secondary wall commences prior to the cessation of the elongation phase and continues to approximately 40 DPA, forming a wall of  
10 almost pure cellulose.

The constructs for use in such cells may include several forms, depending upon the intended use of the construct. Thus, the constructs include vectors, transcriptional cassettes, expression cassettes and plasmids. The transcriptional and  
15 translational initiation region (also sometimes referred to as a "promoter,"), preferably comprises a transcriptional initiation regulatory region and a translational initiation regulatory region of untranslated 5' sequences, "ribosome binding sites," responsible for binding mRNA to ribosomes and translational  
20 initiation. It is preferred that all of the transcriptional and translational functional elements of the initiation control region are derived from or obtainable from the same gene. In some embodiments, the promoter will be modified by the addition of sequences, such as enhancers, or deletions of nonessential and/or  
25 undesired sequences. By "obtainable" is intended a promoter having a DNA sequence sufficiently similar to that of a native promoter to provide for the desired specificity of transcription

of a DNA sequence of interest. It includes natural and synthetic sequences as well as sequences which may be a combination of synthetic and natural sequences.

5 Cotton fiber transcriptional initiation regions chosen for cotton fiber modification may include the 4-4, *rac13* and *Ltp* cotton fiber promoter regions provided herein.

A transcriptional cassette for transcription of a nucleotide sequence of interest in cotton fiber will include in the direction of transcription, the cotton fiber transcriptional initiation  
10 region, a DNA sequence of interest, and a transcriptional termination region functional in the plant cell. When the cassette provides for the transcription and translation of a DNA sequence of interest it is considered an expression cassette. One or more introns may be also be present.

15 Other sequences may also be present, including those encoding transit peptides and secretory leader sequences as desired.

Fiber-tissue transcription initiation regions of this invention are, preferably, not readily detectable in other plant tissues. Transcription initiation regions capable of initiating  
20 transcription in other plant tissues and/or at other stages of fiber development, in addition to the foregoing, are acceptable insofar as such regions provide a significant expression level in cotton fiber at the defined periods of interest and do not negatively interfere with the plant as a whole, and, in  
25 particular, do not interfere with the development of fiber and/or fiber-related parts.

Downstream from, and under the regulatory control of, the cotton fiber transcriptional/translational initiation control region is a nucleotide sequence of interest which provides for modification of the phenotype of fiber. The nucleotide sequence  
5 may be any open reading frame encoding a polypeptide of interest, for example, an enzyme, or a sequence complementary to a genomic sequence, where the genomic sequence may be an open reading frame, an intron, a noncoding leader sequence, or any other sequence where the complementary sequence inhibits transcription, messenger  
10 RNA processing, for example, splicing, or translation. The nucleotide sequences of this invention may be synthetic, naturally derived, or combinations thereof. Depending upon the nature of the DNA sequence of interest, it may be desirable to synthesize the sequence with plant preferred codons. The plant preferred  
15 codons may be determined from the codons of highest frequency in the proteins expressed in the largest amount in the particular plant species of interest. Phenotypic modification can be achieved by modulating production either of an endogenous transcription or translation product, for example as to the  
20 amount, relative distribution, or the like, or an exogenous transcription or translation product, for example to provide for a novel function or products in a transgenic host cell or tissue. Of particular interest are DNA sequences encoding expression products associated with the development of plant fiber, including  
25 genes involved in metabolism of cytokinins, auxins, ethylene, abscissic acid, and the like. Methods and compositions for modulating cytokinin expression are described in United States

Patent No. 5,177,307, which disclosure is hereby incorporated by reference. Alternatively, various genes, from sources including other eukaryotic or prokaryotic cells, including bacteria, such as those from *Agrobacterium tumefaciens* T-DNA auxin and cytokinin biosynthetic gene products, for example, and mammals, for example interferons, may be used.

Other phenotypic modifications include modification of the color of cotton fibers. Of interest are genes involved in production of melanin and genes involved in the production of indigo. Melanins are dark brown pigments found in animals, plants and microorganisms, any of which may serve as a source for sequences for insertion into the constructs of the present invention. Specific examples include the tyrosinase gene which can be cloned from *Streptomyces antibioticus*. The ORF438 encoded protein in *S. antibioticus* also is necessary for melanin production, and may provide a copper donor function. In addition, a tyrosinase gene can be isolated from any organism which makes melanin. The gene can be isolated from human hair, melanocytes or melanomas, cuttle fish and red roosters, among others. See, for example, EP Application No. 89118346.9 which discloses a process for producing melanins, their precursors and derivatives in microorganisms. Also, See, Bernan et al. Gene (1985) 37:101-110; and della-Cioppa et al. Bio/Technology (1990) 8:634-638.

Indigo may be obtained by use of genes encoding a mono-oxygenase such as xylene oxygenase which oxidizes toluene and xylene to (methyl) benzyl alcohol and also transforms indole to indigo. Cloning of the xylene oxygenase gene and the nucleotide



and amino acid sequences are described in unexamined Japanese Patent Application Kokai:2-119777, published May 7, 1990. A dioxxygenase such as naphthalene dioxxygenase which also converts indole to indigo finds use; the naphthalene dioxxygenase gene nahA is described in Science (1983) 222: 167. For cloning, nucleotide sequence in characterization of genes encoding naphthalene dioxxygenase of *Pseudomonas putida*. See, Kurkela et al. Gene (1988) 73:355-362. A tryptophanase gene sequence can be used in conjunction with an oxygenase to increase the amount of indole available for conversion to indigo. Sources of tryptophanase gene sequences include *E. coli* (see, for example, Deeley et al. (1982) *J. Bacteriol.* 151 :942-951).

Plastid targeting sequences (transit peptides) are available from a number of plant nuclear-encoded plastid proteins, such as the small subunit (SSU) of ribulose biphosphate carboxylase, plant fatty acid biosynthesis related genes including acyl carrier protein (ACP), stearyl-ACP desaturase,  $\beta$ -ketoacyl-ACP synthase and acyl-ACP thioesterase, or LHCP II genes. The encoding sequence for a transit peptide which provides for transport to plastids may include all or a portion of the encoding sequence for a particular transit peptide, and may also contain portions of the mature protein encoding sequence associated with a particular transit peptide. There are numerous examples in the art of transit peptides which may be used to deliver a target protein into a plastid organelle. The particular transit peptide encoding sequence used in the instant invention is not critical, as long as delivery to the plastid is obtained.

As an alternative to using transit peptides to target pigment synthesis proteins to plastid organelles, the desired constructs may be used to transform the plastid genome directly. In this instance, promoters capable of providing for transcription of genes in plant plastids are desired. Of particular interest is the use of a T7 promoter to provide for high levels of transcription. Since plastids do not contain an appropriate polymerase for transcription from the T7 promoter, T7 polymerase may be expressed from a nuclear construct and targeted to plastids using transit peptides as described above. (See McBride *et al.* (1994) *Proc. Nat. Acad. Sci.* 91:7301-7305; see also copending US patent application entitled "Controlled Expression of Transgenic Constructs in Plant Plastids", serial no. 08/472,719, filed June 6, 1995, and copending US patent application SN 08/167,638, filed December 14, 1993 and PCT/US94/14574 filed December 12, 1994.) Tissue specific or developmentally regulated promoters may be useful for expression of the T7 polymerase in order to limit expression to the appropriate tissue or stage of development.

Targeting of melanin synthesis genes to vacuoles is also of interest in plant tissues which accumulate the tyrosine substrate involved in melanin synthesis in vacuoles. The protein signal for targeting to vacuoles may be provided from a plant gene which is normally transported across the rough endoplasmic reticulum, such as the 32 amino acid N-terminal region of the metallocarboxypeptidase inhibitor gene from tomato (Martineau *et al.* (1991) *Mol. Gen. Genet.* 228 :281-286). In addition to the signal sequence, vacuolar targeting constructs also encode a

vacuolar localization signal (VLS) positioned at the carboxy terminus of the encoded protein. Appropriate signal sequences and VLS regions may be obtained from various other plant genes and may be similarly used in the constructs of this invention. Numerous  
5 vacuolar targetting peptides are known to the art, as are reviewed in Chrispeels *et al.*, *Cell* (1992) 68:613-616.

The Maize A1 gene which encodes a dihydroflavonol reductase, an enzyme of the anthocyanin pigmentation pathway is one such gene. In cells that express the A1 gene, dihydrokempferol is  
10 converted to 2-8 alkylleucopelargonidin, which may be further metabolized to pelargonidin pigment by endogenous plant enzymes. Other anthocyanin or flavonoid type pigments may also be of interest for modification of cotton cell fibers, and have been suggested for use in plant flowers (for a review of plant flower  
15 color, see van Tunen *et al.*, *Plant Biotechnology Series*, Volume 2 (1990) Developmental Regulation of Plant Gene Expression, D. Grierson ed.). Anthocyanin is produced by a progression of steps from cellular phenylalanine pools. The R and C1 genes are maize regulatory proteins which are active by positively affecting  
20 upstream steps in the anthocyanin biosynthesis from these pools. The R gene is described in Perot and Cone (1989) *Nucl. Acids Res.*, 17:8003, and the C1 gene is described in Paz-Ares *et al* (1987) *EMBO*, 6:3553-3558. Lloyd *et al.* (1992) *Science*, 258:1773-1775 discussed both genes.

25 Although cotton fibers in commercially grown varieties are primarily white in color, other naturally occurring cotton varieties have brown or reddish-brown fibers. Additionally, a

cotton line containing green colored fibers has been identified. Cotton lines providing such fibers are available from various sources, including the BC variety cottons (BC Cotton Inc., Box 8656, Bakersfield, CA 93389) and Fox Fibre cottons (Natural Cotton Colors, Inc., P.O. Box 791, Wasco, CA 93280).

The existence of such colored cotton lines suggests that the precursors required for the anthocyanin pigment pathways are present in cotton fibers cells, thus allowing further color phenotype modifications. Thus, the maize R and C1 genes could be used in enhancing the levels of of anthocyanin produced in fiber cells. As the R and C1 proteins are proteins with a positive control at the regulatory level on anthocyanin pigment precursor biosynthesis, these proteins are expressed in the nucleus, and not targetted to plastids or vacuoles.

For some applications, it is of interest to modify other aspects of the fiber. For example, it is of interest to modify various aspects of cotton fibers, such as strength or texture of a fiber. Thus, the appropriate gene may be inserted in the constructs of the invention, including genes for PHB biosynthesis (see, Peoples et al. *J. Biol. Chem.* (1989) 264: 15298-15303 and *Ibid.* 15293-15397; Saxena, *Plant Molecular Biology* (1990) 15:673-683, which discloses cloning and sequencing of the cellulose synthase catalytic subunit gene; and Bowen et al. *PNAS* (1992) 89:519-523 which discloses chitin synthase genes of *Saccharomyces cerevisiae* and *Candida albicans*. Various constructs and methods are disclosed for the use of hormones to effect changes to fiber quality in copending US patent application entitled "Cotton

Modification Using Ovary-Tissue Transcriptional factors", serial no. 08/397,652 filed February 2, 1995, the teachings of which are incorporated herein by reference.

Transcriptional cassettes may be used when the transcription  
5 of an anti-sense sequence is desired. When the expression of a polypeptide is desired, expression cassettes providing for transcription and translation of the DNA sequence of interest will be used. Various changes are of interest; these changes may include modulation (increase or decrease) of formation of  
10 particular saccharides, hormones, enzymes, or other biological parameters. These also include modifying the composition of the final fiber that is changing the ratio and/or amounts of water, solids, fiber or sugars. Other phenotypic properties of interest for modification include response to stress, organisms,  
15 herbicides, brushing, growth regulators, and the like. These results can be achieved by providing for reduction of expression of one or more endogenous products, particularly an enzyme or cofactor, either by producing a transcription product which is complementary (anti-sense) to the transcription product of a  
20 native gene, so as to inhibit the maturation and/or expression of the transcription product, or by providing for expression of a gene, either endogenous or exogenous, to be associated with the development of a plant fiber.

The termination region which is employed in the expression  
25 cassette will be primarily one of convenience, since the termination regions appear to be relatively interchangeable. The termination region may be native with the transcriptional

initiation region, may be native with the DNA sequence of interest, may be derived from another source. The termination region may be naturally occurring, or wholly or partially synthetic. Convenient termination regions are available from the

5 Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. In some embodiments, it may be desired to use the 3' termination region native to the cotton fiber transcription initiation region used in a particular construct.

- 10 As described herein, in some instances additional nucleotide sequences will be present in the constructs to provide for targeting of a particular gene product to specific cellular locations. For example, where coding sequences for synthesis of aromatic colored pigments are used in a construct, particularly
- 15 coding sequences for enzymes which have as their substrates aromatic compounds such tyrosine and indole, it is preferable to include sequences which provide for delivery of the enzyme into plastids, such as an SSU transit peptide sequence. Also, for synthesis of pigments derived from tyrosine, such as melanin,
- 20 targeting to the vacuole may provide for enhanced color modifications.

For melanin production, the tyrosinase and ORF438 genes from *Streptomyces antibioticus* (Berman et al. (1985) 37:101-110) are provided in cotton fiber cells for expression from a 4-4 and Rac13

25 promoter. In *Streptomyces*, the ORF438 and tyrosinase proteins are expressed from the same promoter region. For expression from constructs in a transgenic plant genome, the coding regions may be

provided under the regulatory control of separate promoter regions. The promoter regions may be the same or different for the two genes. Alternatively, coordinate expression of the two genes from a single plant promoter may be desired. Constructs for  
5 expression of the tyrosinase and ORF438 gene products from 4-4 and rac promoter regions are described in detail in the following examples. Additional promoters may also be desired, for example plant viral promoters, such as CaMV 35S, can be used for constitutive expression of one of the desired gene products, with  
10 the other gene product being expressed in cotton fiber tissues from the 4-4 and rac promoter.

Similarly, other constitutive promoters may also be useful in certain applications, for example the mas, Mac or DoubleMac, promoters described in United States Patent No. 5,106,739 and by  
15 Comai et al., *Plant Mol. Biol.* (1990) 15:373-381). When plants comprising multiple gene constructs are desired, for example plants expressing the melanin genes, ORF438 and tyrosinase, the plants may be obtained by co-transformation with both constructs, or by transformation with individual constructs followed by plant  
20 breeding methods to obtain plants expressing both of the desired genes.

A variety of techniques are available and known to those skilled in the art for introduction of constructs into a plant cell host. These techniques include transfection with DNA  
25 employing *A. tumefaciens* or *A. rhizogenes* as the transfecting agent, protoplast fusion, injection, electroporation, particle acceleration, etc. For transformation with *Agrobacterium*,

plasmids can be prepared in *E. coli* which contain DNA homologous with the Ti-plasmid, particularly T-DNA. The plasmid may or may not be capable of replication in *Agrobacterium*, that is, it may or may not have a broad spectrum prokaryotic replication system such as does, for example, pRK290, depending in part upon whether the transcription cassette is to be integrated into the Ti-plasmid or to be retained on an independent plasmid. The *Agrobacterium* host will contain a plasmid having the *vir* genes necessary for transfer of the T-DNA to the plant cell and may or may not have the complete T-DNA. At least the right border and frequently both the right and left borders of the T-DNA of the Ti- or Ri-plasmids will be joined as flanking regions to the transcription construct. The use of T-DNA for transformation of plant cells has received extensive study and is amply described in EPA Serial No. 120,516, Hoeckema, In: The Binary Plant Vector System Offset-drukkerij Kanters B.V., Alblasterdam, 1985, Chapter V, Knauf, et al., Genetic Analysis of Host Range Expression by *Agrobacterium*, In: Molecular Genetics of the Bacteria-Plant Interaction, Puhler, A. ed., Springer-Verlag, NY, 1983, p. 245, and An, et al., *EMBO J.* (1985) 4:277-284.

For infection, particle acceleration and electroporation, a disarmed Ti-plasmid lacking particularly the tumor genes found in the T-DNA region) may be introduced into the plant cell. By means of a helper plasmid, the construct may be transferred to the *A. tumefaciens* and the resulting transfected organism used for transfecting a plant cell; explants may be cultivated with transformed *A. tumefaciens* or *A. rhizogenes* to allow for transfer



of the transcription cassette to the plant cells. Alternatively, to enhance integration into the plant genome, terminal repeats of transposons may be used as borders in conjunction with a transposase. In this situation, expression of the transposase should be inducible, so that once the transcription construct is integrated into the genome, it should be relatively stably integrated. Transgenic plant cells are then placed in an appropriate selective medium for selection of transgenic cells which are then grown to callus, shoots grown and plantlets generated from the shoot by growing in rooting medium.

To confirm the presence of the transgenes in transgenic cells and plants, a Southern blot analysis can be performed using methods known to those skilled in the art. Expression products of the transgenes can be detected in any of a variety of ways, depending upon the nature of the product, and include immune assay, enzyme assay or visual inspection, for example to detect pigment formation in the appropriate plant part or cells. Once transgenic plants have been obtained, they may be grown to produce fiber having the desired phenotype. The fibers may be harvested, and/or the seed collected. The seed may serve as a source for growing additional plants having the desired characteristics. The terms transgenic plants and transgenic cells include plants and cells derived from either transgenic plants or transgenic cells.

The various sequences provided herein may be used as

25 molecular probes for the isolation of other sequences which may be useful in the present invention, for example, to obtain related transcriptional initiation regions from the same or different

plant sources. Related transcriptional initiation regions obtainable from the sequences provided in this invention will show at least about 60% homology, and more preferred regions will demonstrate an even greater percentage of homology with the probes. Of particular importance is the ability to obtain related transcription initiation control regions having the timing and tissue parameters described herein. For example, using the probe 4-4 and rac, at least 7 additional clones, have been identified, but not further characterized. Thus, by employing the techniques described in this application, and other techniques known in the art (such as Maniatis, et al., *Molecular Cloning, - A Laboratory Manual* (Cold Spring Harbor, New York) 1982), other transcription initiation regions capable of directing cotton fiber transcription as described in this invention may be determined. The constructs can also be used in conjunction with plant regeneration systems to obtain plant cells and plants; thus, the constructs may be used to modify the phenotype of fiber cells, to provide cotton fibers which are colored as the result of genetic engineering to heretofor unavailable hues and/or intensities.

Various varieties and lines of cotton may find use in the described methods. Cultivated cotton species include *Gossypium hirsutum* and *G. babadense* (extra-long staple, or Pima cotton), which evolved in the New World, and the Old World crops *G. herbaceum* and *G. arboreum*.

Color phenotypes can be assessed by the use of a colorimeter, an instrument which is already used to provide objective measurements of the color of cotton samples. A colorimeter uses a

combination of light sources and filters to make various estimates of a samples colors, sometimes referred to as tristimulus values.

In the past such estimates have been used to calculate a value (Hunter's + b, described below) indicating the degree of  
5 yellowness of a cotton sample. The yellowness and reflectance (from Rd, the degree of lightness or darkness of the samples) has been used to provide cotton color measurements for grading. Tests are typically conducted by exposing the face of a sample to a controlled light source. A typical color chart showing how the  
10 official grade standards relate to Rd and+ b measurements is shown in Cotton, RJ Kohel and CF Lewis, Editors #24 in AGRONOMY Series-American Soc. Agromony (see Fig. 12-6).

Various colorimeter methods can be so used to quantify color and express it numerically. The Munsell method, devised by the  
15 American artist A.. Munsell, uses a classification system of paper color chips assorted according to their hue (Munsell Hue), lightness (Munsell Value), and saturation (Munsell Chroma) for visual comparison with a specimen color.

Other methods for expressing color numerically have been  
20 developed by an international organization concerned with light and color, the Commission Internationale de l'Eclairage (CIE), having a Central Bureau located at Kegelgasse 27, A-1030 Vienna, AUSTRIA. The two most widely known of these methods are the Yxy color space, devised in 1931 based on the tristimulus value XYZ,  
25 as defined by CIE, and the L\*a\*b\* color space, devised in 1976 to provide more uniform color differences in relation to visual differences. Color spaces\* such as these are now used throughout

the world for color communication. The Hunter Lab color space was developed in 1948 by R.S. Hunter as a uniform color space which could be read directly from a photoelectric colorimeter (tristimulus method).

- 5        The  $L^*C^*h$  color space uses the same diagram as the  $L^*a^*b^*$  color space, but uses cylindrical coordinates instead of rectangular coordinates. In this color space,  $L^*$  indicates lightness and is the same as the  $L^*$  of the  $L^*a^*b^*$  color space,  $C^*$  is chroma, and  $h$  is the hue angle. The value of chroma  $C$  is 0 at  
10    the center and increases according to the distance from the center. Hue angle is defined as starting at the  $+a$  axis of the  $L^*a^*b^*$  space, and is expressed in degrees in a counterclockwise rotation. Thus, relative to the  $L^*a^*b^*$  space,  $0^\circ$  and  $360^\circ$  would be at the  $+a^*$  line,  $90^\circ$  would be  $+b^*$ ,  $180^\circ$  would be  $-a^*$  and  $270^\circ$   
15    would be  $-b^*$ .

All of the above methods can be used to obtain precise measurements of a cotton fiber color phenotype.

#### EXPERIMENTAL

- 20        The following examples are offered by way of illustration and not by limitation.

##### Example 1

##### CDNA libraries

##### Tissue preparation for cDNA synthesis

- 25        Leaf and root tissue were isolated from 8 inch tall greenhouse grown seedlings and immediately frozen in liquid nitrogen. Flowers were collected at the rapidly expanding 3 day

preanthesis stage and also frozen. Seed was collected from 21 day postanthesis locules which had been removed from the boll and frozen entire in liquid nitrogen. Once frozen, the fiber was removed from the seed and the denuded seed used for RNA isolation.

- 5 All fibers were removed from the seed under liquid nitrogen and the fiber was ground to a powder prior to RNA isolation. Fibers were from bolls which had been tagged at anthesis.

#### DNA and RNA Manipulations

- 10 The lambda ZapII™ cDNA library system of Stratagene was used for screening, and was prepared from cDNA derived from poly-A<sup>+</sup> mRNA isolated from fibers of *Gossypium hirsutum* cultivar Acala SJ-2. The fibers were isolated from bolls harvested at approximately 21 dpa using field-grown plants in Israel.
- 15 Total RNA was isolated from 21 dpa seeds (*G. hirsutum* cv Coker 130 from which the fiber had been removed) using the method of Hughes and Galau ((1988) Plant Mol Biol Reporter, 6:253-257.) All other RNAs were prepared according to Hall et al. ((1978), Proc Natl Acad Sci USA 75: 3196-3200), with the following
- 20 modifications. After the second 2M LiCl wash, the pellet was dissolved in 1/10 original volume of 10 mM Tris pH7.5 and brought to 35mM potassium acetate pH6.5 and 1/2 volume EtOH was added slowly. The mixture was placed on ice for 15 minutes and then centrifuged at 20,000 x g for 15 minutes at 4°C. The potassium
- 25 acetate concentration was brought to 0.2M, 2 1/2 volumes EtOH added and the RNA placed at -20°C for several hours. The precipitate was centrifuged at 12,000 x g for 30 minutes at 4°C

and the pellet was resuspended in diethylpyrocarbonate-treated water. Poly-A<sup>+</sup> RNA was prepared from total mRNA utilizing an oligo(dT)-cellulose kit (Becton Dickenson) and following the manufacturer's protocol.

- 5 Cotton genomic DNA was prepared as follows. Four grams of young cotton leaf tissue (cv Coker 130) was ground to a powder in N<sub>2</sub> and placed in an Oak Ridge tube with 0.4g polyvinylpyrrolidone and 20mls extraction buffer (200mM Ches/NaOH pH9.1, 200mM NaCl, 100mMEDTA/NaOH pH9.0, 2% SDS, 0.5% Na deoxycholate, 2% Nonidet NP-  
10 40, 20mM B-mercaptoethanol) was added to sample, gently mixed and incubated at 65°C in a shaking water bath for 10 minutes. 7.0 mls of 5M potassium acetate pH6.5 was added and carefully mixed. Incubation was carried out on ice for 30 minutes with gentle mixing every 5 minutes. The sample was centrifuged for 20 minutes  
15 at 21,000 x g and the supernatant was filtered through Miracloth into another tube and centrifuged as before. The supernatant was again filtered through Miracloth into 15 mls of room temperature isopropanol in an Oak Ridge tube. After gentle mixing, the sample was incubated at room temperature for 10-60 minutes until the DNA  
20 precipitated. The DNA was spooled and allowed to air dry before being resuspended in 4 mls of TE on ice for 1 hour. CsCl was added to 0.97g/ml final concentration and 300 ul 10mg/ml ethidium bromide was also added before filling VTi80 quick seal tubes. The sample was centrifuged overnight at 225,000 x g overnight. The  
25 DNA was extracted with water saturated butanol and enough water was added to bring the volume to 4 mls before adding 2 volumes

EtOH. The DNA was spooled, air dried and resuspended in 200 ul sterile water.

#### Northern and Southern Analysis

5 For Northern, 10ug of total RNA was isolated from various tissues, separated by electrophoresis in 1.2% agarose-formaldehyde gels and transferred onto Nytran Plus membranes (Schleicher and Schuell). Hybridization conditions consisted of a solution containing 50% formamide(v/v), 5xSSC, 0.1% SDS, 5mM EDTA, 10x  
10 Denhardt's solution, 25mM sodium phosphate pH6.5 and 250 ug/ml carrier DNA. Washes were performed in 2xSSC, 0.1% SDS at 42°C 3 times for 30 minutes each time.

Cotton genomic DNA (12ug) was digested with various restriction endonucleases, electrophoresed in 0.9% agarose gels  
15 and blotted onto Nytran Plus membranes. Hybridization and filter washing conditions for both the 3' specific and full-length cDNA insert probes were as described for Northern analysis.

Probes derived from 3'-untranslated regions were synthesized via oligonucleotide primers from the Rac13 cDNA, corresponding to  
20 bases 600-619 and 843-864 (Figure 4). Each set of primers was used in a polymerase chain reaction to synthesize copies of 3'-specific DNA sequences. These sequences were used as templates in the generation of single-stranded, <sup>32</sup>P-labeled probes off the antisense strand in a polymerase chain reaction. The full-length  
25 cDNA inserts for Rac13 were used as templates for double stranded, random primed probes using the Prime-It kit (Stratagene).

## Example 2

Isolation of cDNA Clones from Cotton

cDNA to the 4-4 clone was isolated from the cotton fiber library described above, and shown to express in fiber but not other tissues. This sequence was not related to any known protein. Only 400 kb of encoding sequence was present in this clone, so the library was rescreened using the cDNA to obtain full-length clones. The full-length encoding sequence is provided in Figure 1.

By comparing sequences of random cDNA clones against various sequence data banks via BLAST, a National Center for Biotechnology Information service, a clone, designated #105, was found to have an encoding sequence related to that of a reported lipid transfer protein.

Another clone was sequenced which showed high homology to animal Rac proteins. This clone, designated Rac, was not quite full-length, and the library was re-screened using this initial Rac DNA segment as probe. Of approximately 130,000 primary plaques screened, 56 screened positive; of these, 14 clones were isolated and sequenced. Of these 14 clones, 12 showed identical sequence homology to the original Rac clone and one of these cDNA clones encoded a full length cDNA and received the name Rac13. Figure 4 shows the cDNA sequence encoding the Rac13 gene expressed in cotton fiber.

One other partial-length cDNA clone, designated Rac9, was clearly related, but distinct in DNA and amino acid sequence from Rac13. Re-screening of 150,000 plaques resulted in the isolation



of 36 positive clones of which only two clones corresponded to the Rac9 sequence (both full-length clones), the remainder being Rac13. These results suggest that cotton contains genes for at least two distinct Rac proteins. Based upon the frequency of  
5 clone isolation, Rac13 is relatively highly-expressed and Rac9 less so in cotton fibers at 21 days post-anthesis (dpa), the age at which polyA<sup>+</sup> mRNA was isolated for library construction.

Comparisons of the deduced amino acid sequence of Rac13 with other small G-proteins showed that the cotton Rac proteins are  
10 very closely related to the Rho1 protein sequence deduced from a cDNA clone isolated recently from pea (Yang and Watson, *supra*). After the pea Rho1, mammalian Rac proteins show the highest homology with the cotton Rac proteins. Other proteins of the rho subfamily, such as the yeast CDC42 and human RhoA, are also  
15 clearly related to the cotton Rac genes. By contrast, the other small G-proteins of the Rab/YPT subfamily isolated from plants such as the example shown of the tobacco RAB5 protein, as well as the human Ras proteins, are least homologous to the cotton Rac proteins of all the small G-proteins compared. The cotton and pea  
20 proteins, as well as the mammalian Racs, all have pI's above 9, whereas those of other rho and ras proteins are in the range of 5.0-6.5.

### Example 3

#### 25 Expression of Cotton Fiber Genes in Developing Fibers

Expression of the Rac13 and 4-4 genes was assessed using mRNA prepared from various cotton tissues and from fibers at

- different stages of development. Blots were hybridized with probes derived from untranslated regions of Ltp, Rac13 and 4-4 genes. The gene for Rac13 exhibits highly-enhanced expression in fibers; virtually no detectable mRNA is present in leaves, roots, or flower parts, even under conditions of extended development time. Rac13 expression is detected in seeds at an age that corresponds to the highest expression levels observed in fiber tissue derived from seeds of this same age. The pattern of Rac13 expression in fibers is very dependent upon the developmental stage. Expression is very low during the stage of primary wall synthesis (0-14 dpa, see Meinert and Delmer, 1977), reaches a maximum during the transition to secondary wall synthesis (about 15-18 dpa), and declining during the stage of maximal secondary wall cellulose synthesis (about 24-28 dpa).
- 4-4 mRNA begins to accumulate in fiber cells only at day 17 post anthesis and continues through at least day 35 post anthesis. Levels peak at day 21 and remain high. 4-4 mRNA is not detected in other cotton tissues, and is not detected in fiber tissue before onset at 17 days post anthesis.
- The #105 lipid transfer protein cDNA clone was used as a probe against cotton tissue and in a cotton fiber northern. The northern showed that the cotton fiber Ltp is highly expressed in cotton fiber. The mRNA that codes for this protein is expressed throughout fiber development at extremely high level. Northern blots indicate that this mRNA is expressed at 5 dpa and is continually expressed at a high level at 40 dpa.

## Example 4

Genomic DNA

cDNA for both the 4-4 and Rac13 was used to probe for genomic clones. For both, full length genomic DNA was obtained from a library made using the lambda dash 2 vector from Stratagene™, which was used to construct a genomic DNA library from cotton variety Coker 130 (*Gossypium hirsutum* cv. coker 130), using DNA obtained from germinating seedlings.

The cotton genomic library was probed with a 3'-specific Ltp probe and 6 genomic phage candidates were identified and purified. Figure 7 provides an approximately 2 kb sequence of the Ltp promoter region which is immediately 5' to the Ltp encoding region.

Six genomic phage clones from the cotton genomic library were identified using a 3'-specific probe for the Ltp mRNA. This was done to select the promoter from the Ltp gene that is maximally expressed in cotton fiber from the family of Ltp genes in cotton. The Ltp promoter is active throughout the fiber development period.

20

## Example 5

Preparation of 4-4 Promoter ConstructspCGN5606

The pCGN5606 promoter construct comprises the 4-4 cotton fiber expression cassette in a first version, version I (Figure 2). The sequences from nt1 to 65 and nt 5,494 to 5,547 correspond to fragments of the pBluescriptII polylinker where this cassette

is cloned. Unique restriction enzyme sites present in these regions flanking the cassette allow the cloning of the fiber expression cassette into binary vectors including the pCGN 5138 and 1547 series.

5       The sequences from nt57 to 5,494 are contained in a lambda phage clone of a cotton Coker 130 genomic library. This lambda genomic clone was given the designation 4-4(6).

      The region from nt 65 to nt 4,163 corresponds to the 5' flanking region of the 4-4(6) gene. At nt 4,163 there is a NcoI  
10       restriction site sequence that corresponds to the first codon of the 4-4 (6)ORF.

      The region from nucleotide 4,163 to 4,502 corresponds to part of the 4-4 (6)ORF. The sequence from nt 4,502 to 4,555 is a  
15       synthetic polylinker oligonucleotide that contains unique target sites for the restriction enzymes EcoRI, SmaI, SalI, NheI and BglII. This fragment from nt4,163 to 4,555 is a stuffer fragment and is left in place to facilitate the monitoring of cloning manipulations.

      The genes to be expressed in cotton fiber cells using this  
20       cassette can be cloned between the NcoI restriction site and any of the polylinker sites. This operation will replace the stuffer fragment with the gene of interest. The region from nt 4,555 to 5,494 corresponds to the 940 nucleotides downstream of the stop codon and constitute the 3' flanking region of the 4-4 (6) gene.  
25       There is a unique AscI restriction enzyme site at nt 5483.

pCGN5610

The pCGN5610 construct is a second version of a 4-4 cotton fiber expression cassette, version II, which is a modified version of pCGN5606. The two versions of the 4-4 cotton fiber expression cassette are designed to allow the cloning of tandem arrays of two  
5 fiber cassettes in one binary plasmid. The differences with respect to pCGN5606 are very minor and described below.

The XbaI restriction site in the region of nt 1 to 65 has been deleted by standard cloning manipulations. The polylinker region is in the reverse orientation of pCGN5606.  
10 There is a unique XbaI restriction enzyme site at nt5484. The sequences from nt1 to 57 and nt 5,494 to 5,518 of pCGN5610 correspond to fragments of the pBluescriptII polylinker where this cassette is cloned. Unique restriction enzyme sites present in these regions allow the cloning of the fiber expression cassette  
15 into binary vectors of the pCGN 5138 and 1547 series.

The sequences from nt57 to 5,494 are contained a lambda phage clone of a Coker 130 genomic library. This clone is described in my notebook as lambda genomic clone 4-4(6). The region from nt 57 to nt 4,155 corresponds to the 5' flanking  
20 region. At nt 4,155 there is a NcoI restriction site sequence that corresponds to the first codon of the 4-4 ORF. The region from nucleotide 4,156 to 4,500 corresponds to part of the 4-4 ORF. This fragment from nt4,156 to 4,550 is a stuffer fragment and is left in place to facilitate the monitoring of  
25 cloning manipulations. The sequence from nt 4,500 to 4,550 is a synthetic polylinker oligonucleotide containing unique target

sites for the restriction enzymes BglII, NheI, SalI, SmaI and EcoRI.

The genes to be expressed in cotton fiber cells using this cassette can be cloned between the NcoI restriction site and any  
5 of the polylinker sites. This operation replaces the stuffer fragment with the gene of interest. The region from nt 4,550 to 5,494 corresponds to the 940 nucleotides downstream of the stop codon and constitute the 3' flanking region of the 4-4 (6) gene.

10

#### Example 6

#### Preparation of Rac13 Promoter Constructs

##### Genomic clone

From a genomic clone designated 15-1, mapping was done with restriction endonucleases. The largest fragment with the Rac13  
15 coding region was identified. This was a Pst fragment, and when subcloned in the Bluescript™ KS+ vector (BSKS+; Stratagene) was named pCGN4722. The insert had a length of 9.2 kb.

The region of the Pst fragment with the Rac13 coding sequence was identified. DNA sequence was determined for approximately 1.7  
20 kb 5' of the start codon and approximately 1.2 kb 3' of the stop codon. The entire Rac coding region (exons and introns) was conveniently flanked by NdeI sites.

pCGN4722 was digested with XbaI, and a 2.7 kb fragment was removed. Religation gave pCGN4730, which was then digested with  
25 NdeI, dropping out a 1.7 kb fragment containing the entire Rac coding region. Religation yielded pCGN4731.

A polylinker region was created using overlapping synthetic oligonucleotides which were PCR'ed using primers homologous to the 5' and 3' ends of the resynthesized section. The resulting product was digested with EcooR1 and Hind III and ligated into BSKS+ at the EcoR1 and Hind III sites. The resulting plasmid was designated pCGN4733.

10 pCGN4731 and pCGN4633 were digested with Nde1 and the Nde1 fragment containing the synthesized polylinker region from pCGN4733 was dropped in the Nde1 site of 4731, giving pCGN4734. This last plasmid was digested with Sal and Xba, and so was pCGN5133. pCGN5133 was the 9.2 kb pst fragment in BSKS+ where the polylinker sites flanking the insert were altered to different sites for ease of manipulation. The fragment from pCGN4734 was then placed into the equivalent site of pCGN5143, giving pCGN4735.

15 A sequence for approximately 3 kb of the promoter construct pCGN4735 is provided in Figure 5. The resynthesized sequence falls between the Nde1 sites located at bases 1706 and 1898 of the sequences. Thus, the sequence in Figure 5 includes approximately 1.7 kb 5' to the Nde1 site 5' to the resynthesized polylinker region. There is a roughly 2.5 kb sequence 5' from this sequence which is not provided in Figure 5, relative to the total 9.2 kb insert. The sequence of Figure 5 also includes approximately 1.1 kb 3' to the 3' Nde1 site. Approximately 3 kb which is most 3' in the Rac13 insert is not provided in Figure 5. A map for pCGN4735 is provided in Figure 6.

#### Example 7

Pigment Synthesis GenesMelanin

A binary construct for plant transformation to express genes for melanin synthesis is prepared as follows. The melanin genes  
5 were originally isolated from the common soil bacterium *Streptomyces antibioticus* (Bernan et al. (1985) 34:101-110). Melanin production is composed of a two gene system. The first gene, *tyrA*, encodes the catalytic unit responsible for the polymerization of the amino acid tyrosine, the primary substrate,  
10 and is termed tyrosinase. The second gene, ORF438, is responsible for binding copper and delivering copper to the tyrosinase and activating the enzyme. Expression of both the ORF438 and *tyrA* genes ensures maximal tyrosinase activity.

The genes for both ORF438 and *tyrA* were fully re-synthesized  
15 with respect to their DNA sequence. This was performed as the initial DNA sequence isolated from *Streptomyces* has a very high guanine and cytosine (G+C) DNA content. Thus, the ORF438 and *tyrA* genes were re-synthesized to appear more "plant-like" (reduced G+C content) with respect to plant preferred codons encoding their  
20 corresponding amino acids.

Indigo

Indigo production involves conversion of the amino acid tryptophan, the primary substrate, into indole which is then  
25 converted into indoxyl. Molecules of indoxyl spontaneously convert to indigo in the presence of oxygen. A two gene system was used to affect indigo production in fiber cells. The first



gene (*tna*) was obtained from the bacterium *E. coli* and encodes the enzyme tryptophanase. The designation *tna* stands for the gene encoding tryptophanase from *E. coli*, an enzyme which converts tryptophan to indole (Stewart et al., (1986) *J Bacteriol* 166:217-223).

The *pig* designation is used for the encoding sequence to the protein for indigo production from *Rhodococcus*, which produces indigo from indole (Hart et al., (1990) *J Gen Microbiol* 136:1357-1363). Both *tna* and *pig* were obtained by PCR. Tryptophanase is responsible for the conversion of tryptophan to indole, while the second gene (*pig*) encodes an indole oxygenase enzyme responsible for the conversion of indole to indoxyl. Both these bacterial genes were utilized in their native form.

#### Example 8

##### Constructs for Targeting Pigment Synthesis Genes

For plastid targeting, the constructs contain a fragment of the tobacco ribulose biphosphate carboxylase small subunit gene encoding the transit peptide and 12 amino acids of the mature protein (Tssu) positioned in reading frame with the appropriate encoding sequence.

For vacuolar targeting of the melanin synthesis genes, constructs include a fragment of the metallocarboxypeptidase inhibitor gene, encoding the entire 32 amino acid N-terminus signal peptide of that protein plus 6 amino acids of the mature protein (CPI+6) (Martineau et al., *supra*), positioned in reading frame with the appropriate encoding sequences. In addition to the

signal peptide, a sequence encoding a vacuolar localization signal (VLS) is inserted 3' of the protein encoding sequence.

Constructs which contain encoding sequences for bacterial genes involved in biosynthesis of pigmented compounds and  
5 sequences for directing transport of the encoded proteins into plastids or vacuoles are prepared as follows.

#### Melanin

The re-synthesized ORF438 and tyrA genes were treated in two  
10 distinct ways depending on which compartment in the fiber cell the final protein products would be localized. One chimeric gene/plant binary construct (designated pCGN5148) contained the genes targeted to the fiber cell plastids. To do this, 12 amino acids of a gene for the small subunit of carboxylase (SSU) plus  
15 the original 54 amino acid SSU transit peptide were fused to the amino termini of both the ORF438 and tyrA gene products respectively. These peptide sequences allow the ORF438 and tyrA gene products (proteins) to be efficiently targeted to the plastid. This targeting was initiated as the plastid is the site  
20 of tyrosine production within the fiber cell.

The second chimeric gene/plant binary construct (designated pCGN5149) contained the ORF438 and tyrA genes targeted to the vacuole within the fiber cell. Based on information from other biological systems, it was postulated that the fiber cell vacuole  
25 may contain a high concentration of tyrosine for melanin polymerization. Both the ORF438 and tryA genes contain the 29 amino acid signal peptide from a tomato carboxypeptidase inhibitor

(CPI) protein as amino terminal gene fusions to direct these proteins to the endoplasmic reticulum (ER) secretory system of the fiber cell.

In addition, the *tyrA* gene has an 8 amino acid vacuolar targeting peptide (VTP) from CPI fused at the carboxy terminus so that the mature copper-activated tyrosinase will eventually be targeted to the vacuole of the fiber cell. Both the ORF438 and *tyrA* proteins also had potential glycosylation sites removed via site-directed mutagenesis of the ORF438 and *tyrA* genes respectively. Potential plant cell glycosylation of these proteins upon their expression in fiber cells could result in tyrosinase inactivation, hence removal of potential glycosylation sites was deemed necessary.

#### 15 Indigo

The only modification to the indigo genes was the fusion of the tobacco SSU transit peptide encoding DNA sequences onto the amino terminal region of both the *tna* and *pig* genes to affect the localization of both the tryptophanase and indole oxygenase proteins to the fiber cell plastid. These are the same exact gene fusions that were made for the plastid-directed proteins for melanin production in construct 5148. The *tna* and *pig* gene products were targeted to the fiber cell plastid as that is the primary site of tryptophan synthesis.

25

#### Example 9

#### Expression Constructs

### Melanin

The modified genes for both the plastid and vacuolar targeted ORF438 and tyrosinase proteins were placed into a fiber expression cassette to be "switched" on during development of the cotton fiber cell. The "switch" (promoter) utilized for the melanin constructs was 4-4. The modified ORF438 and *tyrA* genes were cloned into the 4-4 promoter cassette and these chimeric genes then inserted into a binary plasmid to create plasmids pCGN5148 and pCGN5149, containing the modified genes for plastid and vacuolar targeted ORF438 and tyrosinase proteins, respectively. These binary plasmids also contain genetic determinants for their stable maintenance in *E. coli* and *Agrobacterium* and also contain a chimeric gene for plant cell expression of the bacterial kanamycin resistance gene. This kanamycin resistance marker allows for the selection of transformed versus non-transformed cotton cells when plant hypocotyl or leaf segments are infected with *Agrobacterium* containing the binary plasmids.

A block diagram of the plasmid pCGN5149, having vacuolar targetting sequences, is shown in Figure 8. Plasmid pCGN5148 (not shown) is constructed the same as 5149, only pCGN5148 has plastid-targetting sequences.

### Indigo

As with the melanin genes, the plastid-directed *tna* and *pig* genes were placed in the fiber-specific 4-4 promoter cassette and these chimeric genes subsequently inserted into a binary plasmid

to create plasmid pCGN5616. A block diagram of plasmid pCGN5616 is shown in Figure 8.

#### Anthocyanin

5 A construct has been prepared for the expression of the maize R and CI genes in developing cotton fiber. These genes are known to be responsible for the production of Anthocyanin pigments by acting in a regulatory manner to turn on the chalcone pathway for production of anthocyanins (red spectrum colors). The R and CI  
10 genes were placed under the control of the Rac13 promoter cassette. A binary plasmid designated pCGN4745 (not shown), contains both the R and CI genes each under control of the Rac13 promoter.

15

#### Example 10

##### Cotton Transformation

##### Explant Preparation

Coker 315 seeds are surface disinfected by placing in 50% Clorox (2.5% sodium hypochlorite solution) for 20 minutes and  
20 rinsing 3 times in sterile distilled water. Following surface sterilization, seeds are germinated in 25 x 150 sterile tubes containing 25 mls 1/2 x MS salts; 1/2 x B5 vitamins; 1.5% glucose; 0.3% gelrite. Seedlings are germinated in the dark at 28°C for 7 days. On the seventh day seedlings are placed in the light at  
25 28±2°C.

##### Cocultivation and Plant Regeneration

Single colonies of *A. tumefaciens* strain 2760 containing binary plasmids pCGN2917 and pCGN2926 are transferred to 5 ml of MG/L broth and grown overnight at 30°C. Bacteria cultures are diluted to  $1 \times 10^8$  cells/ml with MG/L just prior to cocultivation.

5 Hypocotyls are excised from eight day old seedlings, cut into 0.5-0.7 cm sections and placed onto tobacco feeder plates (Horsch et al. 1985). Feeder plates are prepared one day before use by plating 1.0 ml tobacco suspension culture onto a petri plate containing Callus Initiation Medium CIM without antibiotics (MS

10 salts: B5 vitamins: 3 % glucose: 0.1 mg/L 2,4-D: 0.1 mg/L kinetin: 0.3% gelrite, pH adjusted to 5.8 prior to autoclaving). A sterile filter paper disc (Whatman #1) was placed on top of the feeder cells prior to use. After all sections are prepared, each section

15 paper towels and returned to the tobacco feeder plates.

Following two days of cocultivation on the feeder plates, hypocotyl sections are placed on fresh Callus Initiation Medium containing 75 mg/L kanamycin and 500 mg/L carbenicillin. Tissue was incubated at 28±2°C, 30uE 16:8 light:dark period for 4 weeks.

20 At four weeks the entire explant was transferred to fresh callus initiation medium containing antibiotics. After two weeks on the second pass, the callus was removed from the explants and split between Callus Initiation Medium and Regeneration Medium (MS salts: 40mM KNO<sub>3</sub>: 10 mM NH<sub>4</sub>Cl:B5 vitamins:3% glucose:0.3% gelrite:400 mg/L carb:75 mg/L kanamycin).

25

Embryogenic callus was identified 2-6 months following initiation and was subcultured onto fresh regeneration medium.

Embryos are selected for germination, placed in static liquid Embryo Pulsing Medium (Stewart and Hsu medium: 0.01 mg/l NAA: 0.01 mg/L kinetin: 0.2 mg/L GA3) and incubated overnight at 30°C. The embryos are blotted on paper towels and placed into Magenta boxes  
5 containing 40 mls of Stewart and Hsu medium solidified with Gelrite. Germinating embryos are maintained at  $28 \pm 2^\circ\text{C}$  50  $\mu\text{E m}^{-2}\text{s}^{-1}$  16:8 photoperiod. Rooted plantlets are transferred to soil and established in the greenhouse.

Cotton growth conditions in growth chambers are as follows:  
10 16 hour photoperiod, temperature of approximately 80-85°, light intensity of approximately 500  $\mu\text{Einsteins}$ . Cotton growth conditions in greenhouses are as follows: 14-16 hour photoperiod with light intensity of at least 400  $\mu\text{Einsteins}$ , day temperature 90-95°F, night temperature 70-75°F, relative humidity to  
15 approximately 80%.

#### Plant Analysis

Flowers from greenhouse grown T1 plants are tagged at anthesis in the greenhouse. Squares (cotton flower buds),  
20 flowers, bolls etc. are harvested from these plants at various stages of development and assayed for enzyme activity. GUS fluorometric and histochemical assays are performed on hand cut sections as described in co-pending application filed for Martineau et al., *supra*. For fiber color characteristics, plants  
25 are visually inspected, or northern or western analysis can be performed, if necessary.

Example 11Expression of Transgenic Pigment Synthesis GenesMelanin

5       Plants that exhibited resistance to the kanamycin selectable  
marker via a leaf assay and corresponding Western analysis were  
considered transformed. Transgenic fiber was collected from  
individual plant transformants at different stages of fiber  
development and analyze in two ways. One was to analyze fiber at  
10 a single developmental time point for each transgenic cotton plant  
to compare tyrosinase expression between transgenic events. The  
other was to screen developing fiber from selected plants to  
analyze the timing of tyrosinase expression under the control of  
the fiber-specific 4-4 promoter, by Western blots using antisera  
15 prepared against purified tyrosinase protein.

For the plastid-targeted construct pCGN5148 9 of 13 events  
screened for tyrosinase expression were positive, while 13 of the  
16 transformed vacuolar-targeted construct pCGN5149 events which  
were screened were positive. Expression level in the fiber in  
20 tyrosinase positive plants is approximately 0.1-0.5% fiber cell  
protein. Clearly, the cotton fiber cells comprising the DNA color  
constructs DNA produce the necessary proteins required for  
synthesis of a pigment.

Visually, the lint from the tyrosinase positive events  
25 exhibits color to varying degrees, while plants that do not  
express the enzyme do not exhibit any color. Colorimeter  
measurements of cotton fiber taken from control Coker 130 plants



and plants from various events transformed with pCGN5148 are provided in Figures 9 and 10, respectively.

Fiber from pCGN5148 (plastid-directed) plants demonstrates a bluish-green color phenotype. One event, 5148-50-2-1 included  
5 cotton fiber cells (linters) which were colored and which had an negative  $a^*$  value less than - 8.0, as measured on the  $L^*a^*b^*$  color space. Coker 130 cotton fiber cells do not typically demonstrate a negative  $a^*$  value.

These colored cotton cells also had a color located on the  
10  $L^*C^*h$  color space with a relatively high hue angle value  $h$ , greater than  $135^\circ$ . Normal Coker 130 fibers have a similar value which is not greater than about  $90^\circ$  as measured by this method.

Results of colorimeter measurements of cotton fiber taken from plants transformed with pCGN5149 are provided in Figure 11.  
15 Fiber from plants expressing tyrosinase from construct pCGN5149 (vacuolar-targetted) tends to have a light brown phenotype.

### Indigo

Resistance to the kanamycin selectable marker via leaf assay  
20 and Western analysis was again the criterion for designating a plant as transformed by pCGN5616. Transgenic fiber was collected from individual plant transformants at different stages of fiber development. The transgenic developing fiber is screened from selected plants to analyze the timing of *tna* and *pig* gene  
25 expression under the control of the fiber-specific 4-4 promoter and fiber is also analyzed at a single developmental time point for each transgenic cotton plant for comparison of both

tryptophanase and indole oxygenase expression between transgenic events, by using Western blots with antisera prepared against the tryptophanase and indole oxygenase proteins.

For the indigo events, 15 of 24 screened plants were positive for expression of both the tryptophanase and indole oxygenase enzymes. Expression levels in the fiber of these proteins is between 0.05-0.5% fiber cell protein. Approximately half of these transformants are expressing both genes in the fiber resulting in a very faint light blue color phenotype. Visually, there is a faint blue color in the majority of these positive events, particularly in 20-30 dpa fiber in the unopened boll. Results of colorimeter measurements of cotton fiber taken from various events of plants transformed with pCGN5616 are provided in Figure 12. Many of these events had relatively low  $a^*$  values (less than 2) with elevated  $b^*$  values (greater than 10), as measured on the  $L^*a^*b^*$  color space. Similarly, several 5149 events also measured with an  $a^*$  value less than 2 while maintaining a  $b^*$  value greater than 10.

#### 20 BC Cotton

Colorimeter measurements taken on naturally colored fiber from four separate BC cotton lines is provided in Figure 13.

The above results demonstrate that the color phenotype of a transgenic cotton fiber cell can be altered by expressing pigment synthesis genes. The transgenic cotton fiber cells include both a pigment synthesizing protein, and pigment produced by the pigment

synthesizing protein. As shown from the results of Figures 9 through 13, expression of a pigment gene of interest can result in cotton fiber cells in which the synthesis of pigments combined with appropriate targeting sequences results in modification of color phenotype in the selected plant tissue, yielding colored cotton fiber by expression from a genetically engineered construct.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application are specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail, by way of illustration and example for purposes of clarity and understanding, it will be readily apparent to those of ordinary skill in the art that certain changes and modifications may be made thereto, without departing from the spirit or scope of the appended claims.

**CLAIMS**

What is claimed is:

1. A DNA construct comprising as operably joined  
5 components in the direction of transcription, a cotton fiber  
transcriptional factor and an open reading frame encoding a  
protein of interest, wherein said transcriptional factor is  
selected from the group consisting of the Ltp, the 4-4 and  
the rac promoter sequences.
- 10 2. The DNA construct according to Claim 1, further  
comprising a transport signal encoding sequence from a plant  
nuclear-encoded gene.
3. The DNA construct according to Claim 2, wherein said  
transport signal encoding sequence comprises a plastid  
15 transit peptid.
4. The DNA construct according to Claim 1, wherein said  
transport signal encoding sequence encodes a signal peptide  
which provides for transport across the rough endoplasmic  
reticulum.
- 20 5. The DNA construct according to Claim 4, wherein said  
sequence further comprises, 3' to said open reading frame, a  
vacuolar localization signal.
6. The DNA construct of Claim 1 wherein said pigment is  
melanin or indigo.
- 25 7. The DNA construct of Claim 6 wherein said open  
reading frame is from a bacterial gene.

8. The DNA construct of Claim 7 wherein said bacterial gene is selected from the group consisting of ORF438, *tyrA*, anthocyanin R gene, anthocyanin C1 gene, *pig*, and *tna*.

9. A plant cell comprising a DNA construct of Claim 1.

5 10. A cotton plant cell according to Claim 9.

11. A cotton fiber cell according to Claim 10.

12. A plant comprising a cell of any one of Claims 9-11.

13. A method of modifying fiber phenotype in a cotton plant, said method comprising:

transforming a plant cell with DNA comprising a construct for expression of a protein in a pigment biosynthesis pathway, wherein said construct comprises as operably joined components:

15 a transcriptional initiation region functional in cells of said cotton plant,

an open reading frame encoding a protein of interest, and

a transcriptional termination region functional in cells of said cotton plant,

20 wherein said plant cell comprises a substrate of said protein; and

growing said plant cell to produce a cotton plant, wherein said protein reacts with said substrate to produce  
25 said pigment.

14. The method of Claim 13 wherein said construct further comprises a transport signal encoding sequence from a plant nuclear-encoded gene.

15 15. The method of Claim 13 wherein said transport signal encoding sequence encodes a signal peptide which provides for transport across the rough endoplasmic reticulum.

16. The method of Claim 13 wherein said DNA comprises constructs for expression of two proteins in a pigment biosynthesis pathway, wherein each of said constructs  
10 comprises components i) through iv), and wherein said two proteins are not encoded by the same gene.

17. The method of Claim 16 wherein said pigment is melanin and said proteins are encoded by *tyrA* and *ORF438*.

15 18. The method of Claim 16 wherein said pigment is indigo and said proteins are *tna* and *pig*.

19. The method of Claim 16 wherein said pigment is anthocyanin and said constructs comprise the anthocyanin R and C1 regulatory genes.

20 20. The method of Claim 13 wherein plant cell is a cotton fiber cell, and wherein said transcriptional region is a fiber tissue transcription initiation region.

21. The method of Claim 20 wherein said transcriptional region is selected from the group consisting of the *Ltp*, the  
25 4-4 and the *rac* promoter sequences

22. A recombinant DNA construct comprising the cotton tissue transcriptional sequence shown in Figure 2.

23. A recombinant DNA construct comprising the cotton tissue transcriptional sequence shown in Figure 5.

24. An isolated DNA encoding sequence of Figure 1.

25. An isolated DNA encoding sequence of Figure 4.

5 26. The method of Claim 13 wherein said protein of interest is involved in the synthesis of a plant hormone.

27. An isolated DNA sequence comprising the cotton lipid transfer protein encoding sequence of Figure 7.

28. A cotton fiber cell comprising a DNA sequence, wherein  
10 said DNA sequence comprises as operably joined components in the direction of transcription, a cotton fiber transcriptional factor and an open reading frame encoding a protein required for synthesis of a pigment.

29. A cotton fiber cell according to Claim 27 comprising  
15 pigment produced by said pigment synthesizing protein.

30. A cotton fiber cell according to Claim 27 wherein said DNA sequence further comprises a transport signal encoding a sequence from a plant nuclear-encoded gene.

31. A cotton fiber cell according to Claim 29, wherein said  
20 transport signal encoding sequence comprises a plastid transit peptid.

32. A cotton fiber cell according to Claim 29, wherein said transport signal encoding sequence encodes a signal peptide which provides for transport across the rough endoplasmic reticulum.

25 33. A cotton fiber cell according to Claim 31, wherein said sequence further comprises, 3' to said open reading frame, a vacuolar localization signal.

34. A cotton fiber cell according to Claim 27 wherein said transcriptional factor is selected from the group consisting of the cotton fiber lipid transfer promoter sequence, the 4-4 promoter sequence and the *rac* promoter sequence.

5 35. A cotton fiber cell according to Claim 27 wherein said pigment is melanin or indigo.

36. A cotton fiber cell according to Claim 27 wherein said open reading frame is from a bacterial gene.

37. A cotton fiber cell according to Claim 35 wherein said  
10 bacterial gene is selected from the group consisting of ORF438, *tyrA*, anthocyanin R gene, anthocyanin C1 gene, *pig*, and *tna*.

38. A cotton fiber cell comprising melanin.

39. A cotton fiber cell comprising indigo.

40. A cotton fiber cell which is colored by genetic  
15 engineering and which has a negative  $a^*$  value less than - 1.0 as measured on the  $L^*a^*b^*$  color space.

41. The cotton fiber cell of Claim 39 wherein said negative  $a^*$  value is less than a -5.0.

42. The cotton fiber cell of Claim 40 wherein said negative  
20  $a^*$  value is less than a -8.0.

43. A cotton fiber cell which is colored by genetic engineering and which has an  $a^*$  value less than 2 and the  $b^*$  value greater than 10 as measured on the  $L^*a^*b^*$  color space.

44. A cotton fiber cell which is colored by genetic  
25 engineering and which has a hue angle value  $h$  of greater than 100° as measured on the  $L^*C^*h$  color space.



45. The cotton fiber cell of Claim 43 wherein said h value is greater than a 135'.

20  
 CTT TCT ATT TGG TTA ACC ATG GCT CAT AAC TTT CGT CAT CCT TTC TTC  
 Leu Ser Ile Trp Leu Thr Met Ala His Asn Phe Arg His Pro Phe Phe>  
 40  
 60  
 CTT TTC CAA CTT TTA CTC ATT ACT GTC TCA CTA ATG ATC GGT AGC CAC  
 Leu Phe Gln Leu Leu Ile Thr Val Ser Leu Met Ile Gly Ser His>  
 80  
 100  
 ACC GTC TCG TCA CGG GCT CGA CAT TTA TTC CAC ACA CAA ACC TCA  
 Thr Val Ser Ser Ala Ala Arg His Leu Phe His Thr Gln Thr Thr Ser>  
 120  
 140  
 160  
 TCA GAG CTG CCA CAA TTG GCT TCA AAA TAC GAA AGG CAC GAA GAG TCT  
 Ser Glu Leu Pro Gln Leu Ala Ser Lys Tyr Glu Lys His Glu Ser>  
 180  
 200  
 220  
 240  
 GAA TAC AAA CAG CCA AAA TAT CAT GAA GAG TAC CCA AAA CAT GAG AAG  
 Glu Tyr Lys Gln Pro Lys Tyr His Glu Tyr Pro Lys His Glu Lys>  
 260  
 280  
 CCT GAA ATG TAC AAG GAG GAA AAA CAA AAA CCC TGC AAA CAT CAT GAA  
 Pro Glu Met Tyr Lys Glu Glu Lys Gln Lys Pro Cys Lys His His Glu>  
 300  
 320  
 GAG TAC CAC GAG TCA CGC GAA TCG AAG GAG CAC GAA GAG TAC GAT AAA  
 Glu Tyr His Glu Ser Arg Glu Ser Lys Glu His Glu Tyr Asp Lys>  
 340  
 360  
 380  
 GAA AAA CCC GAT TTC CCC AAA TGG GAA AAG CCT AAA GAG CAC GAG AAA  
 Glu Lys Pro Asp Phe Pro Lys Trp Glu Lys Pro Lys Glu His Glu Lys>  
 400  
 420

FIGURE 1A

CAC GAA GTC GAA TAT CCG AAA ATA CCC GAG TAC AAG GAC AAA CAA GAT  
 His Glu Val Glu Tyr Pro Lys Ile Pro Glu Tyr Lys Asp Lys Glu Asp>  
 440 460 480  
 GAG AAT AAG AAA CAT AAA GAT GAA GAG TGC CAG GAG TCA CAC GAA TCG  
 Glu Asn Lys Lys Lys His Lys Asp Glu Glu Cys Glu Ser His Glu Ser>  
 500 520  
 AAA GAG CAC GAA GAG TAC GAG AAA GAA AAA CCC GAT TTC CCC AAA TGG  
 Lys Glu His Glu Glu Tyr Glu Lys Glu Lys Pro Asp Phe Pro Lys Trp>  
 540 560  
 GAA AAG CCT AAA GCG CAC GAG AAA CAT AAA GCC GAA TAT CCG AAA ATA  
 Glu Lys Pro Lys Gly His Glu Lys His Lys Ala Glu Tyr Pro Lys Ile>  
 580 600 620  
 CCT GAG TGC AAG GAA AAA CTA GAT GAG GAT AAG GAA CAT AAA CAT GAG  
 Pro Glu Cys Lys Glu Lys Leu Asp Glu Asp Lys Glu His Lys Glu>  
 640 660  
 TTC CCA AAG CAT GAA AAA GAA GAG AAG AAA CCT GAG AAA GGC ATA  
 Phe Pro Lys His Glu Lys Glu Glu Lys Lys Lys Pro Glu Lys Gly Ile>  
 680 700 720  
 GTA CCC TGA GTG GGT TAA AAT GCC TGA ATG GCC GAA GTC CAT GTT TAC  
 Val Pro \*\*\* Val Gly \*\*\* Asn Ala \*\*\* Met Ala Glu Val His Val Tyr>  
 740 760  
 TCA GTC TGG CTC GAG CAC TAA GCC TTA AGC CAT ATG ACA CTG GTG CAT  
 Ser Val Trp Leu Glu His \*\*\* Ala Leu Ser His Met Thr Leu Val His>  
 780 800 \*

FIGURE 1B

```

GTG CCA TCA TCG AGT AAT TTC ATG GGA TAT TGT AAT TAT ATT GTT
Val Pro Ser Cys Ser Asn Phe Met Gly Tyr Cys Asn Tyr Ile Val>

      820
AAT AAA AAA GAT GGT GAG TGG GAA ATG TGT GTG TGC ATT CAT CCA TGA
Asn Lys Lys Asp Gly Glu Trp Glu Met Cys Val Cys Ile His Pro ***>

      840
      860
      880
      900
GCA ATG CTG AAT CTC TTT GCA TGC ATA GAG ATT CTG AAT GGT TAT AGT
Ala Met Leu Asn Leu Phe Ala Cys Ile Glu Ile Leu Asn Gly Tyr Ser>

      920
      940
      960
TTA TGT TAT ATC GTT TGT TCT AGT GAA ATT AAT TTT GAA TGT TGT ATG
Leu Cys Tyr Ile Val Cys Ser Ser Glu Ile Asn Phe Glu Cys Cys Met>

TAA TGT T
*** Cys Xxx>

```

FIGURE 1C

20	40	60
ACTAAAGGGA	ACAAAAGCTG	GAGTCCACC
	GCGTGGCGG	CCGCTCTAGA
		ACTAGTGGAT
80	100	120
	*	
CCCCCGTGGG	CTAAACAAAA	CATGGGAGA
	TTTCTCTTAA	AAAAATMAAA
		GAAGCTTACT
140	160	180
CAATACACT	TTGTGAATTG	TATACAAAG
	ACTCAATGAA	AAACAATAAC
		TCAATACACT
200	220	240
*		
TTTTTTCAC	GATTTACATC	CTTTATATAG
	GCTGAACATA	CAACAACTTT
		AGCTAAAAAA
260	280	300
		*
ATAGGATAAC	CTAATAGCAA	AATCAATC
	AGATATTAAA	CCATGATTTT
		AGCTAACCT
320	340	360
TTAACAACTT	TATTGAAACT	AATTTGAATA
	TTTCACTCTC	TGATATGCC
		AAGATTTTAG
380	400	420
	*	
GCCACTAAC	GAATTTGGGG	TGAACTTTAA
	CATCTCATGC	ATTGTTAAC
		GTTTCAAAACA
440	460	480
AGTTTTTTGC	ATTATTTTAC	TATATGAAC
	GTTTGATTAG	GTTGACTTAC
		ACACTGAGCT
500	520	540
*		
TGTAAGCTCA	CTCAAATTTT	TCTAATTTCT
	AAGGTGATCA	GCAAACTTAG
		GACCGGGGGG
560	580	600
		*
CGTACGAGAG	CTCGGATTGA	TTTTCTAGTT
	AATTAATNAG	ACGATTTATG
		TTTTTAACT

Figure 2a

620	640	660
ATTATGGACT TTTTGGACTA TGTAACGTGT TGGGACTTTA TTTTGTGTTT TTATTGCTT		
680	700	720
TTTTGGATT TAGTAATTAT TATTTTAAA CTGCAAAATT ATATGTTTTT ACAAACTAAG		
740	760	780
TCACAGTTTT CAAAATCCCA TAACTTAGAA TTTTGGCTG CAAATTAAG TAATCAITTA		
800	820	840
AGTGTTTTTT CTGTAATAAA AATAATAAAT AATTTTAAG AGTATTTTCC TAAAAATTGG		
860	880	900
AAATTGATTT ACCAAAATTA GTATGTCAA ACACATGTTT ATATGTTACA GGGCGATATC		
920	940	960
GTCTAGGCAA ATACATCTA GCGCGGGTTT GGAGTGTTAC AGGCGGAGTG GGCTCATTTT		
980	1000	1020
GAGTAAGTAT AGTTAGGCC GAGTTTTAGA TTGCAATATC AAGGTCAAAG ATTTTGTAAA		
1040	1060	1080
CTTCGATGAA TCATATGATT GATTGTCCGA TTAACGAAT ATGTTTTTTT CTTTGTGTG		
1100	1120	1140
TGTTTTATCT CGTGTGATAA GTATATAGTA TGTTTTATTC CAATTCCTAT GGCATGTAC		
1160	1180	1200
ATTGTGGCTA TTCTAATTA APTGATTTGT TATTATGAA ATCTGATCA TGTGTTCTAC		
1220	1240	1260

Figure 2B

AAAGCATGGA ATCTCATGCC TACTGCTTTC TCTTAAGAT ACGATTGCAG GTTAAACATG  
 1280 1300 \* 1320  
 CTTACTATTT TGATTTTGTG CTTGCATGCT ATGTACACAT ACATGGGGTT GGGATGATAT  
 1340 1360 1380  
 GGTAAAGGAG AAGTTTGTAC AGTTTAATGA TTTCGACTAT CTGGTGGTTT AACCAATAT  
 1400 \* 1420 1440  
 TTGTTATGGC ATCTTGACTG CGGTATGGT GGCTGGACCG CCCATATCTG TTCTGGAAAT  
 1460 1480 1500 \*  
 TTATCTGTGA CTCTGGTGGC ATTGTCTACA ATTATTTGTT GGTTGTTTTT GGAATGGACGA  
 1520 1540 1560  
 GTCGTGGGGA ACTCTATTTG GTGTGTTGGC GAGTTGGGTA GGAATTTTTC GAATAAAATTT  
 1580 1600 \* 1620  
 TGCATTGTGT TTTTCTGAAA AATATTGSCAT TAACATAATC ATGCATTCTC AATTTTGGTC  
 1640 1660 1680  
 AATTGAACGT TATAAAATTC TCTATGATAT CCTGATCTGT TTATATACAT ATATGTGTTT  
 1700 \* 1720 1740  
 ATGCTTGAGT TAAGTCAAC ATTGAGATTC ATAGCTACCC CAATTATTTA ATCATTTCAG  
 1760 1780 1800 \*  
 GCAATCTGCA GACTTAGGAT TGGATGGCGT TCAGGAGCTT GGATTTGGTTT TCTCACATCA  
 1820 1840 1860  
 TATTTTATTA AATAATTATT AATTAATAAT TATGGACTTT TGGACTGTCT GACTAATTTT

Figure 2C

1880	1900 *	1920
CAGAAATTTTA TTTTGGTTTT GGGTTTGTGTT GAAATTTTATA GATAATATATT TTAAATATTC		
1940	1960	1980
TGCATAATTT TTCTGTATATT TGAATAAGGAT GTTCGAATTT TTTTTCAAAA TTGAACGTT		
2000 *	2020	2040
TAAGAATTTT TACTACTGCA AATTCAGAAAT AAGTGAATTT GTTTTITAGA AAGATTAAT		
2060	2080	2100 *
AAGTTAGTAT TAGGATTTT AGTTTGATTT GGTGGAAGT AAGTATGTT TTGGAACATA		
2120	2140	2160
ATTATTGAC AATAATTAAG TTTTCTAGG AATTAACGGA AATATCTTCT TCTTTTTTGT		
2180	2200 *	2220
AAATTTACTA ATGCARGAC AATCAACGTT TTGGGGAGCA AATAATCTAG CTTTAAGTAG		
2240	2260	2280
TCAGTGTAACT TCACAAAATC TGGTCATAAC TTCTAGGCTG AGTTTGCTGT GCTACAGTAG		
2300 *	2320	2340
TAAGTCTATA GAACTTACC TGACAAAACG AATGACGTC AGGTCGAAT CTACAACTTT		
2360	2380	2400 *
TCCTTTTCT TCAATTAACA TATGTTTGTAT TCAAGTTCCG ATCTATAATA ATTATTACG		
2420	2440	2460
ATTATCAAT TTCAATTACC TTATATCATC CTATTATAAA TATTAAGTCAG TTCATTTGAG		

Figure 2D



2480	2500	2520
TTTTCGRAAG TTCCAAAAA TTTTGAATTT TATTAAATTT ATTCCCTAAA ACCGAATAG	*	
2540	2560	2580
TTATATCTTT CAATTTAAG TTTCAATTTTT CAATCCGATT TCAATTTTCAT CCTTTTATA		
2600	2620	2640
CCTCTATTA TCTATAATTA CATAAATTC AATTTAATTT TGAATATATT ACACTTAGT		
2660	2680	2700
CCCTAAGTTC AAAACTATA ATTTTCACTT TAGAATTTAA TCATTTTTC CACTTAAGCA		
2720	2740	2760
TCAATTTTAA CCGAATGACA CAATTTTCAT GATTAGTTAG ATCAAGCTTT TGAAGCTTCA		
2780	2800	2820
AAACATAAAA ATTACAAAA AAAACAAAC TTAATATCAT TTATCAATTT GAACAACAAA	*	
2840	2860	2880
GCTTGGCCGA ATGCTAAGAG CTTAAAAATG GCTCTCTTTG TTTCTTTTTG TTGCAACGG		
2900	2920	2940
TGGAGAGAAG AGGGAATGA AGATTGACCA TATTTTTTTA TTATGTTTTA ACAATAATA	*	
2960	2980	3000
TTAATAATTT AATCATAATT ATACTTTGGT GAATGTGACA GTGGGGAGAT ACGTTAAGTA		
3020	3040	3060
TTTTAACATT ATACTTTTTG CACGAGTTG GCTGGTCTAC CCAAGAGTGA TCAAGTTTG		
3080	3100	3120

Figure 2E

9/39

AGTCGCCCTC AATGAGCCAA TTTTGGCCCA TAATGGATAA AGGCAATTTG TTTAGTTCAA \*  
 3140 3160 3180  
 CTGCTCAG AATRACTGTA AATGAAAT AAAATAAGGT GGCTGGTCA CACACAAA  
 3200 3220 3240  
 AAAAAACTAA TGTGGGTGG TTGAATTTTA TATTACGGA TGTATATTTA TATTTTAAAA  
 3260 3280 3300  
 TAAAAATTATG TTTATTTATG TCCTATATTT TTGGAGCAT CCATACTATA ATTGCTAAC \*  
 3320 3340 3360  
 ATATATTTAA AATATAGTAA TATAAAGTGT AATTAACITT AAATTACAAG CATATATTTA  
 3380 3400 3420  
 AATTTTGAAT CAATTAATTT TTATTTCTAT TATTTTAATT AATTAGTCT ATTTTTTCAA  
 3440 3460 3480  
 AATTAATTTT AAATCTAAT AAAAATAAT TTTCCTTAAT GTTGAACAA CTCATGTTAT  
 3500 3520 3540  
 ACTTCAAAAT TATAAGTATT ATATTACCT TGAATGTTTA TTTATTAGTA TATTAATCT  
 3560 3580 3600  
 GATTATAATT ATGGTGGAT ACAATCGCTT TCCACTTAAT ATTTTACTA TGNTTTATTA \*  
 3620 3640 3660  
 ATTTTATTTCA ACAATGATA TTTACTTATT AATACATAAT TTATCATTAAT TTTATGGAAA  
 3680 3700 3720  
 \*

Figure 2F

TTGAGACCA GAACATTAA GAGACAAAT TCTATACAA AGACAATTTA GAAAAAATG  
 3740 3760 3780  
 TACTTTTAGG TAAATTTAAG TACTCTTAAC CAACACAAA AATTCAAATC AAATGAATA  
 3800 3820 3840  
 \*  
 AATAAGATAA TATACATAC GGAACATCTT ACTTGTAATC TTACATTCCTC ATATTTTAT  
 3860 3880 3900  
 \*  
 TATGAAAAAT AATCTTATAT TACTCGAACT AATGTTGTC ACAAATTTAT ATCTAAATAA  
 3920 3940 3960  
 AGAAAAACAC TTAATTTTAA TAACATTTTT TCATATATTT GAAAGATTAT ATTTTGTATA  
 3980 4000 4020  
 \*  
 TTTCGTAATA AATATTTGAC ATAGATTGAG CACCTTCCTA ACATATTCCTC ACCATAGTC  
 4040 4060 4080  
 AAGTATGTAG ATGAGAAATT GGTACAAACA ACGTGGGGCC AAATCCACC AAACCATCTC  
 4100 4120  
 \*  
 TCATTTCTCT CTATAAAAGG CTTCGTACAC ATGACAAACA ATCCACACA C AAA TAC  
 <Phe Val  
 4140 4160 4180  
 ACG TTC TTT TCT TAT TTG ATT AAC CAT GGC TCA TAG CAT TCG TCA  
 <Arg Glu Lys Arg Glu Ile Glu Asn Val Met Ala \*\*\* Leu Met Arg \*\*\*  
 4200 4220  
 \*  
 CCC TTT CTT CCT TTT CCA ACT TTT ACT CAT AAG TGT CTC ACT AGT GAC  
 <Gly Lys Lys Arg Lys Trp Ser Lys Ser Met Leu Thr Glu Ser Thr Val

Figure 2G

11/39

4240 CGG TAG CCA CAC TGT TTC GGC AGC GGC TCG ACG TTT ATT CGA GAC ACA 4280  
 <Pro Leu Trp Val Thr Glu Ala Ala Ala Arg Arg Lys Asn Ser Val Cys  
 4300 \* 4320  
 AGC AAC CTC ATC AGA GCT CCC ACA ATT GGC TTC AAA ATA CGA AAG CAC  
 <Ala Val Glu Asp Ser Ser Gly Cys Asn Ala Glu Phe Tyr Ser Leu Val  
 4340  
 GAG AGT CTG AAT ACG AAA ACG CAG AAT ACA AAC AGC CAA AGT ATC ACG  
 <Leu Thr Gln Ile Arg Phe Ala Leu Ile Cys Val Ala Leu Thr Asp Arg  
 4400 \* 4420  
 AAG AGT ACT CAA AAC TTG AGA AGC CTG AAA TGC AAA AGG AGG AAA AAC  
 <Leu Thr Ser Leu Val Gln Ser Ala Gln Phe Ala Phe Pro Pro Phe Val  
 4440 4460  
 AAA AAC CCT GCA AAC AGC ATG AAG AGT ACC AGC AGT CAC ACG AAT CAA  
 <Phe Val Arg Cys Val Ala His Leu Thr Gly Arg Thr Val Arg Ile Leu  
 4480 4500 \* 4520  
 AGG AGC AAA AAG AGT ACG AGA AAG AAA ATC TCGACGAA TTCGCCCGG  
 <Pro Ala Phe Leu Thr Arg Ser Leu Phe Asp  
 4540 4560 4580  
 CGTCGACGGC TAGCGAAGAT CTTGGGGGCC GTCCAGCCTT GAATCATATG ACACGTGTGC  
 4600 \* 4620 4640  
 ATGTGCCATC ATCAGCACT AATTTCTAGG TATATCGTAA TATATAGTTA ATAAAAAGA  
 4660 4680 4700 \*  
 TGGTGATTGG GAAATGTGTG TGTGCATCC TCCATGCACT AATGGTGAAT CTCCTTGCAAT

Figure 2H

4720 ACATAGAAAT TCTAAATGGT TATAGTTTAT GTTATAGTGT ATGTTGTAGT GAAATTAAAT 4760  
 4740  
 4780  
 4800 \*  
 4820  
 4840  
 4860  
 4880  
 4900 \*  
 4920  
 4940  
 4960  
 4980  
 5000 \*  
 5020  
 5040  
 5060  
 5080  
 5100 \*  
 5120  
 5140  
 5160  
 5180  
 5200 \*  
 5220  
 5240  
 5260  
 5280  
 5300 \*  
 5320

TTAATACTTG TATCTAATGT TACATCACT TGGCTTGATT TATGTTAATG TATGTAATTT  
 ACTTTAATGA TATGCAATGT ATGTTAAT TACATCTGT TGATCAATAT ACTCTTCTAC  
 TATTAATTAAT AAATGCCACT GTTTTGTTTA AACTTTTAC AAGTTAAGAC ATGTATTAAT  
 ATATGACAAT ATAAATTACAG GTTTTAGTTC AATGTTAGCT AUCTTAGTAT GTTATGATG  
 ATCTTAATTA CATTAAACA AATCCACTT AAAAATTTAA TAAATAATAA CAAATAATTA  
 TTGTAATATA ATACATTAAA TGCAACAAA AATGAATAA ATAAATATA ATAGCAATA  
 ATTGTTATA TATTGTAATA TAATATGTAC CATATCTTA ACTGAATAG GGTCTAACCT  
 ATAATCCCTA AAATTTCACT TTAAATATTT TTATACCTAC CATATTAATA GAATCTTTT  
 TAAATATATT AAAATTTTAA TTATCCAT TTAATTAAC TATTAATTAAT CTAACTAAA

Figure 2I

5320                    5340                    5360  
 ATCTAAAATT TTATTTRACC TATTATATA TTCTTATTA TCTTATCTAA TTATAAAGTC  
  
 5380                    5400                    5420  
 TAAATTATCCT AATTATTAATT AAATTCCTTAA TTAATCTTAAT TTGTAACCTC CTCACCCAG  
  
 5440                    5460                    5480  
 CTAGATGCTG GACCCGAATC CGGAGATTTA CATCGGCCAT TGAGATGGCG TGATCAGGGT  
  
 5500                    5520                    5540  
 TTGGGGGGCC GGTACCCAT TCGCCCTATA GTGAGTTGGT ATTACGGCG CTCACTGGT  
 CCGGTTT

Figure 2J

14/39

20	40	60
ACTAAGGGA	ACAAAGCTG	GAGTCCACC
	GGGTGGGG	CCGCTCTAGG
		ATCCCCGGTG
80	100	120
	*	
GACTAAACAA	AACATGGGAA	GATTTGCTCT
	AAAAAATAA	AGAGAGCTTA
		CTCAATAACA
140	160	180
CTTTGTGAAT	TGTATACAAA	AGACTCAATG
	AAAAACAATA	ACTCAATACA
		CTTTTTTICA
200	220	240
*		
CTGATTTACA	TCCTTTATAT	AGGCTGAAC
	TACACAACT	TTAGCTAATA
		AAATAGGATA
260	280	300
		*
ACCTAATAGC	AAATACACAA	TCAGATATTA
	AACCATGATT	TTAGCTAACC
		ATTTAACAAC
320	340	360
TTTATTGAAA	CTAATTTGAA	TATTTCACT
	GCTGATATGC	CCAAGATTTT
		AGGCCACTAA
380	400	420
	*	
CCGATTGGT	GGTCACTTT	AACATGTCAT
	GCATTGTAA	CTGTTTGAAA
		CAAGTTTTTT
440	460	480
GCATTATTTT	ACTATATGAA	CTGTTTGATT
	AGGTTGAGTT	ACACACTGAG
		CTTGTAAAGCT
500	520	540
*		
CACCTCAAAAT	TTTCTAATTT	CTAAGTGAT
	CACCAACTT	AGGACCGGGC
		GGCGTAGGAG
560	580	600
		*
AGCTCGGATT	GAATTTCTAG	TTAATAATA
	AGACGATTA	TGTTTTTAAA
		CTATTATGGA

Figure 3A

620 CTTTITGGAC TATCTAACTG TTGGGACTT TATTTTGT TTTTATTGC TTTTITGGA 660  
 680 700 \* 720  
 TTTAGTAATT ATTATTTTAA AACTGCAAA TTTATATGTTT TTACAACTA AGTCACAGTT  
 740 760 780  
 TTCAAATTC CATTACTTAG AATTTTTCG TCCAAAATAA AGTAATCAAT TAAGTGTTTT  
 800 \* 820 840  
 TTCTGTAATA AAATAATAA ATAATTTTAA CGAGTATTTT CCTAAAAANT GGAAATGAT  
 860 880 900 \*  
 TTACCAAAAT TAGATATCTA AAACACATGT TTTATATGTTA CAGGGCGATA TCGTCTAGGC  
 920 940 960  
 AAATACATC TAGCGGGGT TTGGAGTGTT ACAGGGCGAG TGGGCTCAAT TTGAGTAAAT  
 980 1000 \* 1020  
 ATAGTTAGG CCGAGTTTAA GATTGCATAT TCAAGGTCAA AGATTTTGTA AACTTCGATG  
 1040 1060 1080  
 AATGATATGT ATGATTGTC GATTACGAA ATATGTTTTT TCTTTTGTG TGTGTTTTAT  
 1100 \* 1120 1140  
 CTGCTGTGAT AAGTATATAG TATGTTTTAT TCCAAATCTT ATGGCAGTGT ACATTTGGC  
 1160 1180 1200 \*  
 TATTTCTAAT AAATGATTT GTTATATTG AAATCTGATG CATCTGTCTT ACAAGCATG  
 1220 1240 1260

Figure 3B



16/39

GAATCTCATG CCTACTGCTT TCTGTTAAAG ATACGATTGC AAGTTTAACA TGCTTACTAT  
 1280 1300 1320  
 TTTGATTTTG TCCTTGCTATG CTATGTCACA TTACATGGGG TTGGGATGAT ATGATAAGGA  
 1340 1360 1380  
 GGAAGTTTGG ACAGTTTAAAT GATTTCGACT ATCTGTTGGT TTAACACAT ATTTGTTTATG  
 1400 1420 1440  
 GCACTCTTGC TGGCGTTATG GTGGCTCGAC CGCCCATATC TGTCTCTGGA ATTATCTGT  
 1460 1480 1500  
 GACTCTGGTG GCATTGTCTA CAATTATTG TTGGTGTGTT TTGGATGGAC GAGTCGTGGG  
 1520 1540 1560  
 GAACTCTAAT TGGTGTGTTG CGGAGTTGGG TAGGAATTT TCGAAAAAA TTTGCAATTG  
 1580 1600 1620  
 GTTTTCTCGA AAAATATTGC ATTACATAA TCATGCATTC TCAATTTGG TCAATTGAAC  
 1640 1660 1680  
 GTTATAAAAT TCTCTATGAT ATCTGAACT GTTTATTACA TTATATGTTG TTATGCTTGA  
 1700 1720 1740  
 GTTAAGTCAA ACATTGAGAT TCATGCTCA CCCAATTATT TAATCATTC AGGCAATCTG  
 1760 1780 1800  
 CAGACTTAGG ATTGGATGSC GTTCAGGAGC TTGGATTGGT TTTCACAT CATATTTTAT  
 1820 1840 1860  
 TAAATAATTA TTAATTAATA TTTATGGACT TTGGGACTGT CTGACTAATT TTCGAATTT

Figure 3C

1880	1900	1920
TAATTTGGTT TTGGGTTTTG TTGAATTTTT TAGATAATTA TTTTAAATAT TCTGCATNAT		
1940	1960	1980
TTTTCTGTTA TTGGAAGG AGTTCGAAT TTTTTTCNA AATGAAACG TTAAAGAAT		
2000	2020	2040
TTTACTACTG CAAATTCAGA ATAAATGAAT TTGTTTTTTA GAAGATTAATA ATAAGTTAGT		
2060	2080	2100
ATTACGAATTT TTAGTTTGAT TTGGTGAAA GTAATGTAAG TTTTGAACA TAATATTTTG		
2120	2140	2160
ACAAATAATTA AGTTTCTAG GGAATAACG GAATAATCTT CTTCCTTTTT GTAAAAATTAC		
2180	2200	2220
TAATGCACGA ACAACAACG TTTTGGGAG CAAATAATCT AGCTTTAAGT AGTCAGTGTA		
2240	2260	2280
ACTCTCAAA TCTGGTCATA ACTCTAGGC TGAGTTTCT GTGCTACAGT AGTAAGTCTA		
2300	2320	2340
TAGAAACTTA CCTGCACAAA CGCATGACG TCAGGTCGA ATCTACAACT TTTCCTTTTT		
2360	2380	2400
CTTCAATTA CATATGGTTG ATTCAAGTTC CGATCTATAA TAATTTATTA CGATTTATCA		
2420	2440	2460
ATTTCAATTA CCTTATATCA TCTTATATA ATATATAATC AGTTCAATTC AGTTTTCGAA		

Figure 3D

18/39

2480 2500 \* 2520  
 AGTTCCTAAA AATTTTGAAT TTTATTAAT AAACCGAAAT AGTTATATCT  
 2540 2560 2580  
 TTCAAAATTTA AGTTTCATTT TTCAATCCGA TTTCAAATTC ATCCTTTTAT AACTCTCTAT  
 2600 \* 2620 2640  
 TATCTATAAT TACATAAATT TCAAAATTAAT TTGGAATAT TTACACTTTA GTCCCTAAGT  
 2660 2680 2700 \*  
 TCAAAACTAT AAAATTTTAC TTGGAATTTT AATCAATTTT CACATCTAAG CATCAAAATTT  
 2720 2740 2760  
 AACCMAATGA CACAAATTC ATGATTAAGTT AGATCAAGCT TTTGAGTCTT CAAAACATAA  
 2780 2800 \* 2820  
 AAATTACAAA AAAAAACAA ACTTAAATC AATTATCAAT TTGAACAACA AAGCTTGGCC  
 2840 2860 2880  
 GAATGCTAAG AGCTTAAAAA TGGCTTCTTT TGTTCCTTTT TGTTCGAAAC GGTGGAGAGA  
 2900 \* 2920 2940  
 AGAGGGAAAT GAAGATTGAC CATATTTTTT TATTATGTTT TAACATATAA TATTTAATAAT  
 2960 2980 3000 \*  
 TTAATCATAA TTATACTTTG GTGAATGTA CAGTGGGGAG ATACGTTAAG TATTTTAACTA  
 3020 3040 3060  
 TTATACTTTT TGCACAGAT TGGCTGGTCT ACCCAAGAGT GATCAAAAGTT TGAGCTGCCT  
 3080 3100 3120

Figure 3E

TCAATGAGCC AATTTTGGC CATAATGGAT AAAGGCAATT TGTTTAGTTC AACTGTCAC  
 \*  
 3140 3160 3180  
 AGAATAATGT TAAATGAAA TTAAATAAG GTGGCTGGT CACACACACA AAAAAAACT  
 3200 3220 3240  
 AATGTGGTT GGTGCAATTT TATATTACG ANTCTAATAT TATATTTTAA AATTAATTA  
 \*  
 3260 3280 3300  
 TGTATTATTAG ATTCTTAATA TTITGGAGCA TTCCATACTA TAATTTGGTA ACATAATATT  
 \*  
 3320 3340 3360  
 AAAATATAGT AATAAAGT GTAATTAACT TTAAATTACA AGCAATAATAT TAAATTTTGA  
 3380 3400 3420  
 ATCAATTAAT TTTTATTCTT ATTATTTTAA TTAAATTAGT CTATTTTTC AAATTAAT  
 \*  
 3440 3460 3480  
 TTAAATCTAA ATAAAAATAA TTTTTCCTTA ATGTGGAAC AACTCATGTT ATACTCACA  
 3500 3520 3540  
 ATTATAAGTA TTATATTAC CTGATGAGTT TATTTATTAG TATATTAATT CTGATTATTA  
 \*  
 3560 3580 3600  
 TTATGGTGGG ATCAATCGC TTCCACTTAA ATATTTTAACT TATGATTTAT AAATTTATTT  
 3620 3640 3660  
 CAACATCGTA TATTTACTTA TTAATACATA ATTTATCATA ATTTATGGA AATTGAGACC  
 3680 3700 3720  
 \*

Figure 3F

AAGAAACATT AAGAGACAA ATTCTATAC AAGACAAAT TAGAAAAAA TGTACTTTTA  
 3740 3760 3780  
 GGTAATTTTA AGTACTCTTA ACCAAACCA AAAATTCAAA TCMAATGAAC TAAATAGAT  
 3800 3820 3840  
 AATAAATAC ACGGAACATC TTACTTGTTAA TCTTACNTC CCATAATTTT ATTATGAAA  
 3860 3880 3900  
 ATAACTTTAT ATTACTCGAA CTAAATGTTG TCACAAATTA TTATCTAAT AAGAAAAAC  
 3920 3940 3960  
 ACTTAATTTT TATAACATTT TTTCATATAT TTGAAGATT ATATTTTGTA TATTTACGTA  
 3980 4000 4020  
 AAAATATTTG ACATAGATTG AGCACTTCT TAACTATATC CCACCATAG TCAATATGT  
 4040 4060 4080  
 AGATCGAATA TTGGTACAAA CAACGTGGGG CCAATCCCA CCAACCATC TCTCATTTCT  
 4100 4120  
 TCTATRAAA GCGTTGCTAC ACATAGACAA CAATCCACAC A CA AAT ACA CGT TCT  
 4140 4160 4180  
 TTT CTT TCT ATT TGA TTA ACC ATG G CTCATAGCAT TCGTACCCCT TTCTTCCTTT  
 <Lys Lys Arg Asn Ser \*\*\* Gly His  
 4200 4220 4240  
 TCCAACTTTT ACTCATAAAT GTCTCACTAG TGACCGGTAG CCACACTGTT TGGGACGGG  
 4260 4280 4300

Figure 3g

CTGACGCTTT ATTGAGACA CAAGCAACT CATCAGAGCT CCCACAATTG GCTTCAAAT \*  
 4320 4340 4360  
 ACGAAAGCA CGAAGAGTCT GAATACGAA AGCCAGATA CAACAGCCA AAGTATCAG  
 4380 4400 4420  
 AAGAGTACT AAACTTTGAG AAGCTGAAA TGCAAAAGGA GGA AAAACAA AAACCTGCA  
 4440 4460 4480  
 AACAGCATGA AGAGTACCAC GAGTCACAG AATCAAGGA GC AAAAAGAG TACGAGAAG  
 4500 4520 4540  
 AAAATCTGA CGGCGCGGA GATCTTCGCT AGCGTCCGAC GCCCGGGGA ATTGTCGAG  
 4560 4580 4600 \*  
 CCTTGATCA TATGACGCTG GTGCATGTGC CATCATCATG CAGTAATTC ATGGTATATC  
 4620 4640 4660  
 GTAATATATA GTTAATAAAA AAGATGTGA TTGGGAATG TGTGTGTGA TTCTCCCATG  
 4680 4700 4720  
 CACTAATGGT GAATCTCTTT GCATACATAG AAATCTAAA TGGTTATAGT TTATGTTATA  
 4740 4760 4780  
 GTGTATGTG TAGTGAAAT AATTTAAAT GTTGATCTA ATGTTAATAT CACTTGGCTT  
 4800 4820 4840  
 GATTTATGTT ATGTTATGTA TTTTACTTTA ATGATATGTC ATGTATTGTT AATTTAAT  
 4860 4880 4900 \*

Figure 3H

22/39

TGCTGTGATCA TTATACCTCTT CTACATATTA TTATAAATGG CACTGTTTGG TTAAACTTTT  
 4920 4940 4960  
 TTACAAGCTTA AGACATCTAT AAATATATGCA CAATATAATT ACAAGTTTTA GTTCATGTGT  
 4980 5000 5020  
 AGCTATCTTA GTATGTTTAT GATGATCTTA ATTACATTTTA AACAAATCC ACTTAAATTT  
 5040 5060  
 TTAAATTAATA ATACAAATA ATTATTGTAA TATAATACAT TAAATGCAAC AAAAAATGAA  
 5100 5120 5140  
 ATAAATRAAA TAAAATAGCA AATAATTTGT ATAATATTGT AATATAATAT GTACCAATTT  
 5160 5180 5200 \*  
 CTTAACTGAA ATAGGGCTTA ACCTATATATC CCTAAAATTT CAGTTTAAAT ATTTTATATC  
 5220 5240  
 CTGCCATTT ATTAGACTC TTTTAAATA TATTAAAATT TTAATTATATC CAATTTAATT  
 5280 5300 5320  
 TAAACTATTA ATTATCTTAA CTAAATATCTA AAATTTTATT TAACTATTA ATTAATATCC  
 5340 5360  
 TAAATATCTT ATCTTAATTA AAATCTAAT TATCTTAATT TGATTTAAAT TCTTGATAT  
 5400 5420 5440  
 CTTAATTTGT AACCTCTTCC ACCCAGCTAG ATGCTGGACC CGAATCCGGG AGATTACATC  
 5460 5480 5500 \*  
 GGCATTGAGA TGGCCTAGTA GTGATCAGGG TTTTCTAGAG GTACCCAATT CGCCCTATAG

Figure 31

TGAGTCGT

Figure 3J



50  
 AAAAAACA ATG AGC ACT GCA AGA TTT ATC AAG TGT GTC ACG GTC GGT GAT  
 Met Ser Thr Ala Arg Phe Ile Lys Cys Val Thr Val Gly Asp  
 10  
 98  
 GGA GCT GTG GGG AAA ACT TGT ATG CTC ATT TCA TAT ACC AGC AAT ACT  
 Gly Ala Val Gly Lys Thr Cys Met Leu Ile Ser Tyr Thr Ser Asn Thr  
 15 20 25 30  
 146  
 TTC CCA ACG AGT TAT GTT CCA ACA GTA TTT GAT AAC TTT AGT GCC AAT  
 Phe Pro Thr Asp Tyr Val Pro Thr Val Phe Asp Asn Phe Ser Ala Asn  
 35 40 45  
 194  
 GTG GTG GTG GAT GGC ACG ACA GTG AAC CTT GGC CTA TGG GAC ACT GCC  
 Val Val Val Asp Gly Ser Thr Val Asn Leu Gly Leu Trp Asp Thr Ala  
 50 55 60  
 242  
 GGG CAA GAA GAT TAT AAT AGG CTA AGG CCA CTG AGT TAT AGA GGA GCT  
 Gly Gln Glu Asp Tyr Asn Arg Leu Arg Pro Leu Ser Tyr Arg Gly Ala  
 65 70 75  
 290  
 GAT GTG TTT TTG TTG GCC TTT TCT CTT ATA AGC AAG GCC AGT TAT GAA  
 Asp Val Phe Leu Leu Ala Phe Ser Leu Ile Ser Lys Ala Ser Tyr Glu  
 80 85 90  
 338  
 AAC ATC TAC AAA AAG TGG ATC CCA GAG CTA AGA CAT TAT GCT CAT AAT  
 Asn Ile Tyr Lys Lys Trp Ile Pro Glu Leu Arg His Tyr Ala His Asn  
 95 100 105 110  
 386  
 GTA CCA GTT GTG CTT GTT GGA ACC AAA CTA GAT TTG CGA GAT GAC AAG  
 Val Pro Val Val Leu Val Gly Thr Lys Leu Asp Leu Arg Asp Lys  
 115 120 125  
 434  
 CAG TTC CTC ATT AAT GAT CAC CCT GGA GCA ACA CCA ATA TCA ACA TCT CAG  
 Gln Phe Leu Ile Asp His Pro Gly 135  
 130 140  
 482  
 GGA GAA GAA CTA AAG AAG ATG ATA GGA GCA GTT ACT TAT ATA GAA TGC

FIGURE 4A

Gly Glu Glu Leu Lys Lys Met Ile Gly Ala Val Thr Tyr Ile Glu Cys  
 145 150 155  
 AGC TCC AAA ACC CAA CAG AAT GTG AAG GCT GTT TTC GAT GCT GCA ATA 530  
 Ser Ser Lys Thr Gln Gln Asn Val Lys Ala Val Phe Asp Ala Ala Ile 170  
 160 165  
 AAA GTA GCT TTG AGG CCA CCA AAA CCA AAG AGA AAG CCT TGC AAA AGG 578  
 Lys Val Ala Leu Arg Pro Lys Pro Lys Arg Lys Pro Cys Lys Arg 190  
 175 180 185  
 AGA ACA TGT GCT TTC CTT TGAATATTGG ATCATTATTTA CAGTCAAAAA 626  
 Arg Thr Cys Ala Phe Leu 195  
 CAGTTAACAA AAGCTGTTGC AGATRAACAC TGAATCTGCT ATAGTTTGT TTGGTTTAC 686  
 ATATGTTCCA CGTGAACTA TGAAGCATCT CTAGAGAAAC CCAAACTATC ATATCAACCC 746  
 ATCGATCAAT GAATCGAATT CAATTTTCGC AGTATPAGTT CCTTTAAAC CTITCTTTTT 806  
 ACTTCATTTT ATACGAAATT CTATGNTAA TGTTCCTTAC AAACATGTCA TTACAATGTT 866  
 TAATTATAAA TTCCATCTCT CTATTTTACT AAAAAAAA AAAA 910

FIGURE 4B

26/39

	20	40	60
	TTGGATGAGA	ACCAATTTTT	AATAGTAAGN CCTAACCAAT TTTTAATAAT AAGCTGACT
5	80	100	120
		*	
	CCTAGTACAA	GAGCTTTTAT	TCATTCCTCT ATTTGCTTT CCTCTAGGCT TGGCAATCGA
10	140	160	180
	GAAATTTTCTT	GTGTACAAT	ATAATAATA CATGTAGAA ATAAATTTTA TTCAAATTGA
	200	220	240
	*		
	AGCTTAACC	ATCTTTAATA	TTCTAGATG TAATTTAAT GAAAGATAAA TACATACTCT
15	260	280	300
			*
	TGGACATGA	TTTTCACTT	AATGTTTGT GCTTGGTGA TAGGTGTATT GATGTACGAT
	320	340	360
	GTCTTTTAAA	TCACATATCA	CATTTGACT TTGTATGATG ATAGTCGCAC ATTAACGAAA
20	380	400	420
		*	
	TATGGTGGA	TCTTCACTT	TGAATTGA TAAGTCACCA AACTTTACA AAGTTGATT
25	440	460	480
	GTGTACATAT	ATATATATAT	CTTCAATTT TATATATAA ATTGTGTTTA AATAATTAC
30	500	520	540
	*		
	AGTTATATTA	TTTTTTTATC	TCTAATTTTA TTGTGGCCA AATTTTATG TGAATATTTA
	560	580	600
			*
35	ACATAAAAAA	AATGTGTAC	ATTTACAAGC CCATATACAA ATAAATTATAT AAATATTCAT

FIGURE 5/A

27/39

5                   620                   640                   660  
 TAAAAATAT ATTTAAATAT AGGATATAA TATAACTATT TTAGAATAT TCTACTTTAA  
  
                   680                   700                   720  
 GATAACATAG GTTAAATGTA TTAATTATTA GGTAGTTTA TTGTAAGAT GAGTATATAT  
  
                   740                   760                   780  
 GTCTGAACA TAATCACTAA CCATTTTAT TAACCTCTTG GTTTGAAGT TCCAAAAGA  
  
 10                  800                   820                   840  
 AATGGAAGG GAAATTGAG AGTAAGTTCA TGTATTATTT ATACATAATG AAGTTGATGT  
  
                   860                   880                   900  
 TTTCTTCTTT TTAATATTTT TATACAAAT AITTAATTA AATAATTAG GATTGATCA  
  
 15                                   920                   940                   960  
 AAAATATAAT GAAAGTCGTT TTACTAATAG TCATATTGCA TTTTGTGCGA TCTACTTAAA  
  
                   980                   1000                  1020  
 TAATAGATTA ATTAATTGTS GTACATTAGA TCAAGAACA AACTAGATTT TGTCCCATTC  
  
                   1040                   1060                   1080  
 TATTGTTTAA AGCTGGTCCG TTTACATTAA AATAAGGTAC ATGTTACATG CCACGTATAA  
  
                   1100                   1120                   1140  
 CTATCTGGTT ATTCTATCAA TCAGCTAAT TTTTAACAGT AGAAATGAAT GTAATTTTAA  
  
                   1160                   1180                   1200  
 AATGAAAAGG GTCAAAATGT TATTGATCT AACACGTAGG GATTATTTTA CTATTTTCC  
  
                   1220                   1240                   1260

FIGURE 5/B

**FIGURE 5/C**

29/39

1880 1900 \* 1920  
 ATAAACACTG AATCTGCTAT AGTTTGTTTT TGCTTTCAT ATGTTCCACG TGAACCTATG  
 1940 1960 1980  
 AAGCATCTCT AAGAAACCC AACATCATC ATCAACCCAT CGATCAATGA ATCGATTCCA  
 2000 2020 2040  
 ATTTTCCGAG TATAAGTTCC TTTTAAATCT TTCTTTTAC TTCATTTTAT AACGAATCT  
 2060 2080 2100 \*  
 ATGGATAATG TTCCCTACAA ACATGTCATT ACAATGTTA ATTATTAATT CCATCTCTCT  
 2120 2140 2160  
 ATTTTACTAA GATATTAGTA ACTTCAAACT GCTGATTTT ACTAATTTAT TATTATATAA  
 2180 2200 2220  
 TTGTAGAAAT GATTATTTTT CAATAATTTA ACAACAAAT TTAATATTTAT TATTATATT  
 2240 2260 2280  
 ATTCTCAAT TTTTATTAAA CAAAACATA AATTTTGAC AAATTAAAT AAATGAATTA  
 2300 2320 2340  
 ATTTCTCAAT TTTTCGTGCA ACTATTACAA AAATCTTCA TAGTCTTAAT CTTAATTTGA  
 2360 2380 2400 \*  
 TGCAGAGGTG ATAATAATCT TAATTTGATG CAGAGGTAAT AATGGCCGG GTTTGAGCTG  
 2420 2440 2460  
 GACTTAAAGCA TGATATTGAC GTACTTTATA TTTTCCAAA TTCAACCCAG CTCGAATAT

FIGURE 5/D

5                   2480                   2500 \*                   2520  
 GAGCTAATAA TTTTGTCCAA TTTAATCCAA GCCATTTTA AGTCGTCCA TATTATTTTT  
                   2540                   2560                   2580  
 TAAATTAATA AATTATATC ATTTTATTTT AATATTATAT TATTTTATAT ATTTTTTATT  
                   2600                   2620                   2640  
 TATTGAAAT TTTTATATAG TCACTTAAAC ATTAAGTTAA TGTTTATATT AGAGTAGTAT  
                   2660                   2680                   2700 \*  
 TATATATATT TAGTATAGGT TTAATTTGTT AATAAACTTA AAAATGGGTC TTGTGGGCTA  
 15                   2720                   2740                   2760  
 GACTTGGACC TTAAGGCTC AACTCAAC TTAATTCATA TTTTAAACAG GCTTAATATT  
                   2780                   2800                   2820  
 TTTATTTACA CTGTTTCAAA TTTTTCGGGT GAAATATCTT CGAGCTAGA TTAATAACAC  
                   2840                   2860                   2880  
 CACAGGCTA ATTTGATGCT CAATGAAAT GAAATCATAT TGAGCTTAAT TATATTTCCA  
 25                   2900                   2920                   2940  
 TTCTTCTTTG CTGAAGGAC CAAGCAATC GAGTTACATT AAGGTTAAG AGTATGGGAT  
                   2960                   2980                   3000 \*  
 CGGCCAACC TGCCCCAATG TCTCTTCAAC CATCCAAAA CITGAGTCAG TATCACATAC  
 30                   3020                   3040  
 ATGTACCGNT ATTTATTAT TTAATGAAT TGGCATTTAT TCTTGG  
 35

FIGURE 5/E




FIGURE 6



GGGCAATTCCA CACGACCATG TGTCGCCCTAT TTCCAGGCAAT TTGAGAGCTT CACCTAAACT 60  
TCTAGAGTTG TTTCAAAATTA GCCCCTATTT GTTCTTAAT CAITTTAGGA TCTTGTAAC 120  
TCGTATTAG GACTAAATGT GTAATTTATA CTTTAATTAT GATTGATTA TTAGATTGATT 180  
TNGTAGTAAT GCCCGTGACC CTAATCCGTT AGCGAAGAGG GGTAGGGGT TAGGGGTTTT 240  
ATTATATT TTATAGATT GTATACTCT TGTTTTATTT TTAATTTTGT TACTATTCA 300  
AAGCAATTTG TTGTAGTGT TATTTGAGT AGTTTTATG GGTGAACAC CCTTGACCGC 360  
CAATCAATC ACAAGAGTTC AACATTTTAT TTATTTTGA AATGATTA AATCCTTAAT 420  
CTATATATTC GCCCATTAT TGGGATTTAA TATTCACAG GGTTTAGACC GTCATGAGAC 480  
AGATTAGTTT TATCTTACTG ATGTCACAT CACATAGTA ATTCAACTTA ATACGAGAGG 540  
AACCATTGAT TCACGCCAAT GGTCAATGCA CTTAGTTGAA AAGTAGGGG TCGAAGCTA 600  
CCGTACCTG GATTATGATT GAACACCTCT AAGTCAGAT CCGAATTAGA ACAAATGCAC 660  
GTGTCGGTGG CCTGATTTGCC AACCCCAATA ACAGTGTTG TAGGTTTAA CATTGTTATG 720  
AAAGATAAGG TTTTTTTTTT TATAAGCAAG CAACTATAGG GPTTACTTC CGTGGGCAAA 780  
TTTTTAGGTT ACCTATTTTG GGAGGGGGA TTATGATTCA AGTAAAGAA AGTTGGCACA 840  
CACACAATCA GTACATCTGT TTGACACAG ACACGCCTA AAACAGCAG CAAACAAGCC 900  
TAAGGAATC ACCCAAAAC AACACCAAA AGTACAGAG AAAACAAG AATCCCTGTT 960  
ACCACCAAG TGAATAAAG AAAATAAAC TCACTTTTG GCAATATAA CCTCCTACC 1020  
CTCAACCCCT AACCAAGCAA CAATCAGCAA TACTCCAGC AACCATTTTC CTACAGATT 1080

FIGURE 7A

TGTGTTTCTT GTGATTAATC CAT ATG GCT AGC TCC ATG TCC CTT AAG CTT GCA 1133  
 Met Ala Ser Ser Met Ser Leu Lys Leu Ala>  
 TGT CTG CTA GTG TTG TGC ATG GTG GGT GCA CCC CTG GCT CAA GGG 1181  
 Cys Leu Leu Val Leu Cys Met Val Gly Ala Pro Leu Ala Gln Gly>  
 GAC GTA ACC CGT GCT GAT GGC GTA ACC CTT CCA CGC TGC CTT CCT 1229  
 Asp Val Thr Arg Ala Asp Gly Val Val Thr Leu Pro Arg Cys Leu Pro>  
 TTA TTG ATA GGG AAT GGT AAT GGT GCT GAT GCT GAT GTT GAT GCC CCA 1277  
 Leu Leu Ile Gly Asn Gly Asn Gly Ala Asp Val Asp Ala Pro>  
 GCT TGC TGC GAC ATC GTC AGG GGT CTC TTG AGC TCG CTC TGT GGT 1325  
 Ala Cys Cys Asp Ile Val Arg Gly Leu Leu Ser Ser Leu Cys Gly>  
 GGT GTT TAGGAACCG ATCTAGCTTG AAATCGGGTT CGGATACGGG TGGAGTTTCA 1380  
 Gly Val>  
 AATTGGGTG TTATGGAATC CCAACTTAAT CGTGTTTAGG GGTGGGATCC AATTGTGTGA 1440  
 TACATTACAG AGCATGGTTG TGGATTCTTT TCICATATGT TTGATTCAC TTGCTTGATA 1500  
 CAITGGATGA TTGATTAAGG TGACCGGTTT ACCTGGGTAT CCAACCATCA TCCGATTACT 1560  
 TTTTAAATAT TAATTTGTTT TCCTTTATGT TGCTGTCTT TTGTTTCTT GATCTATAAC 1620  
 ATTAATTTTG CCCAAATTTT CGCATTTTCC ATATGTAGCT TATATATGTA TATATATATT 1680  
 CAATTAAGTA TAATGATTTA GCAGATGATT TGTTATATA TTTAAATCAA ATCAACAATT 1740  
 AATGATCAAT CACTAGCGTC TTAATCTTGA AAAATTCATC AACGGTTATC CTTTGCAGCA 1800  
 TATATATAAA AAATTGCCAA CCCTATGCTT TTACACTTAA TTCAAGGGAT AACATAAGTC 1860  
 GATTAATAACG A 1871

FIGURE 7B




FIGURE 8

Coker 130	Ym, Y	Ym, X	Ym, Y	Lab, L	Lab, A	Lab, B	LCh, L	LCh, A	LCh, H
1	90.35	3206	0.3269	91.84	0.16	5.51	91.84	5.51	88.4
2	77.82	3232	0.3282	90.6	0.66	6.45	90.6	8.48	84.2
3	80.98	3197	0.3257	92.12	0.13	5.04	92.12	5.04	88.6
4	80.16	3200	0.3255	91.75	0.35	5.00	91.75	5.01	86.1
5	77.03	3220	0.3271	90.53	0.81	5.84	90.53	5.87	84.1
6	73.67	3258	0.3283	88.78	1.35	7.14	88.78	7.26	79.4
7	82.43	3178	0.3237	92.76	0.15	4.05	92.76	4.05	87.9
8	82.21	3196	0.3255	92.68	0.19	4.99	92.68	4.99	87.9
9	81.19	3194	0.3241	92.21	0.77	4.42	92.21	4.48	80.2
10	76.11	3243	0.329	89.9	0.74	8.89	89.9	6.82	84
11	82.28	3178	0.3238	92.69	0.19	4.00	92.69	4.00	87.3
TOTAL	874.03	3.5302	3.5883	1005.82	5.30	59.33	1005.62	59.61	939.10
MEAN	79.46	3209	3282	91.42	0.48	5.39	91.42	5.42	85.28
S.D.	2.81	.0026	.0020	1.33	0.38	1.08	1.33	1.11	3.22
RANGE	82.43-73.87	3058-3178	0.3293-3286	92.76-88.76	1.35-0.13	7.14-4.00	92.76-88.76	7.26-4.00	88.6-79.4
AVER DEV.	2.44	.0021	.0017	1.11	0.31	0.88	1.11	0.90	2.84
Coker 130	Hunter L	Hunter a	Hunter B						
1	89.83	0.15	5.42						
2	88.10	0.66	6.27						
3	89.98	0.13	4.98						
4	89.53	0.36	4.94						
5	87.76	0.81	5.69						
6	85.83	1.35	6.85						
7	90.79	0.15	4.03						
8	90.67	0.19	4.85						
9	90.10	0.78	4.38						
10	87.23	0.75	6.65						
11	90.70	0.19	3.98						
TOTAL	980.32	5.32	56.14						
MEAN	89.12	0.48	5.29						
S.D.	1.85	0.39	0.89						
RANGE	90.79-85.83	1.35-0.13	6.85-3.98						
AVER DEV.	1.37	0.31	0.81						
				FIGURE 9					

36/39

5148	Yxy, Y	Yxy, x	Yxy, y	Lab, L	Lab, a	Lab, b	LCh, L	LCh, C	LCh, h
68-1	60.76	0.34	0.35	82.24	2.32	15.11	82.24	15.28	81.3
68-1	61.89	0.34	0.34	82.82	1.97	14.31	82.85	14.44	82.2
50-2-1	78.39	0.3324	0.3375	90.95	0.68	11.29	90.95	11.31	86.6
50-2-1 (lint fiber)	21.49	.3155	0.3489	53.48	-8.01	7.97	53.48	11.29	135.2
5148	Hunter L	Hunter a	Hunter B						
68-1	77.94	2.25	13.35						
68-1	78.67	1.92	12.75						
50-2-1	88.53	0.69	10.71						
50-2-1 (lint fiber)	46.35	-6.35	6.06						
				FIGURE 10					

37/39

5149	Yxy, Y	Yxy, x	Yxy, y	Lab, L	Lab, a	Lab, b	LCh, L	LCh, C	LCh, h
68-1	85.75	0.3351	0.34	84.86	0.72	11.9	84.86	11.92	86.8
68-1	82.54	0.3458	0.3474	83.19	2.14	15.84	83.19	15.98	82.4
68-1	82.56	0.3458	0.3474	83.2	2.14	15.85	83.2	15.99	82.4
8-1	84.72	0.3196	0.3278	83.76	0.89	5.97	83.76	5.93	88.8
68-1	84.97	0.3316	0.3354	84.46	1.17	9.81	84.46	9.87	83.3
17-2	64.42	0.3423	0.3438	84.18	2.28	14.19	84.18	14.36	81
17-3	60.97	0.3475	0.3475	82.36	2.74	16.03	82.36	16.26	80.4
17-15-1	64.02	0.3433	0.3444	83.97	2.34	14.57	83.97	14.75	80.9
21-1	59.32	0.3443	0.3445	81.46	2.84	14.41	81.46	14.84	79.7
21-3	63.54	0.34	0.3409	83.77	2.4	12.88	83.77	13.11	79.5
21-6	67.12	0.3372	0.3394	85.56	1.88	12.15	85.56	12.29	81.3
50-3-1	61.26	0.3502	0.3511	82.51	2.4	17.69	82.51	17.79	82.3
67-1	64.34	0.3434	0.3442	84.13	2.48	14.58	84.13	14.78	80.4
68-1	64.12	0.3442	0.3447	84.02	2.69	14.85	84.02	15.07	80.2
68-2	70.21	0.3428	0.3447	87.09	2.05	15.04	87.09	15.17	82.3
68-3	63.81	0.3457	0.3468	83.86	2.95	15.76	83.86	15.93	81.6
5149	Hunter L	Hunter a	Hunter B						
68-1	81.08	0.71	10.89						
68-1	79.08	2.08	14						
68-1	79.09	2.09	14.02						
8-1	92.04	0.91	5.81						
68-1	80.6	1.15	9.06						
17-2	80.25	2.21	12.75						
17-3	78.08	2.68	14.09						
17-15-1	80.01	2.29	13.05						
21-1	77.01	2.56	12.73						
21-3	79.77	2.35	11.65						
21-6	81.92	1.86	11.14						
50-3-1	78.26	2.33	15.38						
67-1	80.2	2.43	13.07						
68-1	80.07	2.53	13.28						
68-2	83.79	2.04	13.66						
68-3	79.87	2.3	14						

FIGURE 11

38/39

5616	Y94. Y	Y94. X	Y94. Y	Lab. L	Lab. a	Lab. b	Lab. L	Lab. a	Lab. b	Lab. L	Lab. a	Lab. b	Lab. L
11-1	72.28	0.3215	0.3264	88.09	1.1	5.06	88.09	5.17	77.8				
11-2	58.69	0.3284	0.3306	81.12	0.8	8.38	81.12	8.38	85.9				
11-2	52.78	0.3358	0.3395	77.74	3.65	9.22	77.74	9.87	69				
11-1	72.03	0.3312	0.3336	97.96	1.72	8.94	97.96	8.94	78.8				
11-1	72.34	0.3295	0.332	88.13	1.79	8.94	88.13	8.94	78.4				
11-1	71.98	0.3285	0.3313	87.95	2.09	8.39	87.95	8.39	78.1				
11-1	73.01	0.3266	0.3305	88.45	0.89	7.51	88.45	7.54	84.9				
17-1-2	75.85	0.3274	0.3303	89.78	1.62	7.94	89.78	8.08	78.3				
17-3-1	72.6	0.3271	0.3303	88.25	1.48	7.66	88.25	7.8	78.1				
17-4-1	69.02	0.3352	0.3377	86.51	1.78	11.37	86.51	11.5	81.2				
25-11-1	69.5	0.3384	0.3401	86.75	1.88	12.41	86.75	12.47	84.2				
25-25-1	72.21	0.3344	0.3343	86.06	2.09	9.9	86.06	10.11	78.2				
25-35-2	70.46	0.3327	0.3353	87.22	1.73	10.22	87.22	10.38	80.5				
35-35-1	75.59	0.3268	0.3299	89.06	1.58	7.58	89.06	7.73	78.4				
50-12-1	73.13	0.3284	0.3316	88.5	1.46	8.36	88.5	8.48	80.1				
KS-11-2	65.33	0.3371	0.3388	84.65	2.07	11.83	84.65	12	80.1				
5616	Hunter L	Hunter a	Hunter B										
11-1	85	1.09	4.89										
11-2	78.61	0.58	7.64										
11-2	72.64	3.38	8.22										
11-1	84.87	1.74	8.97										
11-1	85.05	1.79	8.2										
11-1	84.84	2.08	7.96										
11-1	85.44	0.67	7.18										
17-1-2	87.08	1.52	7.62										
17-3-1	89.2	1.48	7.31										
17-4-1	83.07	1.76	10.52										
25-11-1	83.98	1.25	1.43										
25-25-1	84.97	2.08	9.32										
25-35-2	83.84	1.72	9.56										
35-35-1	88.84	1.57	7.29										
50-12-1	85.51	1.46	7.96										
KS-11-2	80.82	2.04	10.81										
				FIGURE 12									

[illegible]



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C12N 15/29, 15/82, 5/10, A01H 5/00</b>	<b>A3</b>	(11) International Publication Number: <b>WO 96/40924</b> (43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: <b>PCT/US96/09897</b> (22) International Filing Date: <b>7 June 1996 (07.06.96)</b> (30) Priority Data: 08/480,178 7 June 1995 (07.06.95) <b>US</b> (60) Parent Application or Grant (63) Related by Continuation US Filed on Not furnished (CIP) Not furnished (71) Applicant (for all designated States except US): <b>CALGENE, INC. [US/US]; 1920 Fifth Street, Davis, CA 95616 (US).</b> (72) Inventors; and (75) Inventors/Applicants (for US only): <b>McBRIDE, Kevin [US/US]; 1309 Marina Circle, Davis, CA 95616 (US). STALKER, David, M. [US/US]; 2735 Cumberland Place, Davis, CA 95616 (US). PEAR, Julie, R. [US/US]; 818 Douglass Avenue, Davis, CA 95616 (US). PEREZ-GRAU, Luis [ES/US]; 1230 Elk Place, Davis, CA 95616 (US).</b> (74) Agents: <b>SCHWEDLER, Carl, J. et al.; Calgene, Inc., 1920 Fifth Street, Davis, CA 95616 (US).</b>	(81) Designated States: <b>AU, CA, CN, JP, KG, KZ, MX, TJ, TM, TR, US, UZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b> <b>Published</b> <i>With international search report. Before the expiration of the time limits for amending the claims and to be republished in the event of the receipt of amendments.</i> (86) Date of publication of the international search report: 6 February 1997 (06.02.97)	
(54) Title: <b>COTTON FIBER TRANSCRIPTIONAL FACTORS</b>		
(57) Abstract <p>Novel DNA constructs are provided which may be used as molecular probes or inserted into a plant host to provide for modification of transcription of a DNA sequence of interest during various stages of cotton fiber development. The DNA constructs comprise a cotton fiber transcriptional initiation regulatory region associated with a gene which is expressed in cotton fiber. Also provided is novel cotton having a cotton fiber which has a natural color introduced by the expression in the cotton fiber cell, using such a construct, of pigment synthesis genes. Cotton fiber cells having color produced by genetic engineering and cotton cells comprising melanin and indigo pigments are included.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SS	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

# INTERNATIONAL SEARCH REPORT

Int ional Application No  
PCT/US 96/09897

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/29 C12N15/82 C12N5/10 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,95 08914 (AGRACETUS) 6 April 1995 see page 8, line 12 - page 9, line 17 see page 18, line 6 - page 20, line 16 see page 36, line 1 - page 38, line 36 see sequence ID nos 4 and 5	1
X	WO,A,94 12014 (AGRACETUS) 9 June 1994 see page 9, line 29 - page 10, line 32 see page 19, line 5 - page 21, line 24 see page 40, line 8 - page 43, line 35 --- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not to conflict with the application but cited to understand the principle or theory underlying the invention

- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*A\* document member of the same patent family

2

Date of the actual completion of the international search

6 December 1996

Date of mailing of the international search report

20.12.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 345-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Maddox, A

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/09897

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 9021 Derwent Publications Ltd., London, GB; Class A35, AN 98-161489 XP002020420 & JP.A.02 104 773 (NAGASE SANGYO KK) , 17 April 1990	38
A	see abstract	13-21, 27-36
X	--- TRENDS IN BIOTECHNOLOGY, vol. 10, no. 5, 1 May 1992, pages 165-170, XP000272383 JOHN M E ET AL: "GENES FOR JEANS: BIOTECHNOLOGICAL ADVANCES IN COTTON" see figure 2C	38
P,X	--- WO,A,95 16783 (CALGENE INC ;MCBRIDE KEVIN E (US); STALKER DAVID M (US)) 22 June 1995 see page 8, line 38 - page 10, line 14	13-18,20
P,X	--- US,A,5 495 070 (JOHN MALIYAKAL) 27 February 1996 see sequence ID no. 4. see column 27 - column 28; example 3 see column 45 - column 46	23,26
P,X	--- MOLECULAR AND GENERAL GENETICS, vol. 248, July 1995, pages 43-51, XP002018965 DELMER, D.P., ET AL.: "Genes encoding small GTP-binding proteins analogous to mammalian rac are preferentially expressed in developing cotton fibers" see the whole document	24
A	--- EP,A,0 409 629 (CALGENE INC) 23 January 1991 see the whole document	1
A	--- CROP SCI 25 (5). 1985. 792-797. , XP000612251 KOHLE R J: "GENETIC ANALYSIS OF FIBER COLOR VARIANTS IN COTTON GOSSYPIMUM-HIRSUTUM." see the whole document	13-21, 27-36
A	--- TWELFTH ANNUAL CONFERENCE OF THE SOCIETY OF ETHNOBIOLOGY, RIVERSIDE, CALIFORNIA, USA, MARCH 30-APRIL 2, 1989. J ETHNOBIOL 9 (2). 1989. 239. , XP000612252 FOX S V: "NATURALLY OCCURRING COTTON LINT COLORS." see the whole document	13-21, 27-36
	---	

# INTERNATIONAL SEARCH REPORT

Int. Patent Application No.  
PCT/US 96/09897

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHEMISTRY AND INDUSTRY, 5 September 1994, pages 676-679, XP000466067 JOHN, M.E.: "Re-engineering cotton fibre" see page 677, left-hand column, paragraph 2; table 2 ---	13-21, 27-36
A	WO,A,89 08145 (BIOSOURCE GENETICS CORP) 8 September 1989 see page 5, line 8 - line 18 see page 28, line 17 - line 20 see page 41, line 21 - page 42, line 2 see page 49, line 8 - line 17 ---	16,17
A	WO,A,95 13386 (BIOSOURCE TECH INC) 18 May 1995 see page 23, line 26 - page 24, line 35 ---	16,17
A	WO,A,93 02195 (COMMM SCIENT IND RES ORG) 4 February 1993 see page 1, line 23 - line 29 see page 1, line 36 - page 2, line 2 see page 7, line 1 - line 7 ---	16,17
A	EP,A,0 198 288 (ADVANCED GENETIC SCIENCES INC) 22 October 1986 see page 7, line 32 - line 55 ---	18
T	US,A,5 530 185 (MARTINEAU BELINDA M ET AL) 25 June 1996 see column 8, line 23 - column 9, line 14; claims 1-18; example 10 ---	13,14, 16-20
T	WO,A,96 26639 (CALGENE INC) 6 September 1996 see page 20, line 8 - page 21, line 17 see page 47, line 11 - page 49, line 6 -----	13,14, 16,17

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int ional Application No  
PCT/US 96/09897

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9508914	06-04-95	AU-A- 7925094 EP-A- 0670670 JP-T- 8509871	18-04-95 13-09-95 22-10-96
WO-A-9412014	09-06-94	AU-A- 5676394 CA-A- 2127807 EP-A- 0625006	22-06-94 09-06-94 23-11-94
WO-A-9516783	22-06-95	US-A- 5576198 CA-A- 2178645 EP-A- 0734446	19-11-96 22-06-95 02-10-96
US-A-5495070	27-02-96	US-A- 5521078	28-05-96
EP-A-0409629	23-01-91	WO-A- 9101324 US-A- 5175095 US-A- 5530185	07-02-91 29-12-92 25-06-96
WO-A-8908145	08-09-89	AU-B- 638411 AU-A- 4072589 EP-A- 0406267 JP-T- 3502886 US-A- 5529909 US-A- 5316931	01-07-93 22-09-89 09-01-91 04-07-91 25-06-96 31-05-94
WO-A-9513386	18-05-95	US-A- 5529909 AU-A- 1173495 EP-A- 0731843 ZA-A- 9408973	25-06-96 29-05-95 18-09-96 08-02-96
WO-A-9302195	04-02-93	AU-B- 655494 CA-A- 2112998 EP-A- 0599868 JP-T- 7501686 NZ-A- 243594	22-12-94 04-02-93 08-06-94 23-02-95 26-08-94
EP-A-0198288	22-10-86	US-A- 5180873 AU-B- 602306 AU-A- 5615586 US-A- 5278057	19-01-93 11-10-90 23-10-86 11-01-94

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 96/09897

Patent documents cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5530185	25-06-96	US-A-5175095	29-12-92
		EP-A-0409629	23-01-91
		WO-A-9101324	07-02-91
WO-A-9626639	06-09-96	AU-A-5363296	18-09-96

# Indigo production in developing cotton fibers through the expression of tryptophanase and indole oxygenase



locules dissected 30-35dpa



# Pigment production in genetically engineered cotton fiber

